



Immunostimulant and Pharmacological Activities, of *Cucurbita Maxima* Seeds on Humoral and Cellular Immunity Cells and their Functions

Soukaina Laaraj¹, Ibtissam Ouahidi², Nada Al Moudani¹, Mustapha Allali² and Lotfi Aarab^{1,*}

¹Laboratory of Microbial Biotechnology and Bioactive Molecules (LM2BM), Faculty of Sciences and Techniques of Fez, Sidi Mohammed Ben Abdellah University Morocco

²High Institute of Nursing and Technical Health (ISPITS). Ministry of Health, Fez, Morocco

Abstract:

Background: Natural resources are key for discovering medicinal compounds. Hence, the objective of this study was to evaluate the immunomodulatory activities of *Cucurbita maxima* (pumpkin) seeds, a component of Moroccan folk medicine believed to enhance immunity.

Methods: We produced three distinct extracts employing ethanol, ethyl acetate, and water as solvents. The immunomodulatory effects of these extracts were assessed on rabbit immune cell proliferation and their functions, including IgG production, cytotoxicity, and phagocytosis.

Results: The study findings reveal a significant stimulation in thymocyte proliferation, yielding a remarkable 285.7% response, along with a 146.5% increase in cytotoxicity (MLR) in response to the aqueous extract. Furthermore, there was a notable enhancement in complement activity by 119.9% in response to the aqueous extract, implying the potential of *C. maxima* to modulate both humoral and cellular immunity. Additionally, both EtOAc and aqueous extracts led to a reduction in macrophage phagocytic function. The assessment of the antibacterial properties of ethanol extract showed greater effectiveness against *Bacillus*, *E. Coli*, and *Staphylococcus*. In a separate analysis, antioxidant activity was gauged through DPPH, FRAP, and ABTS methods. These tests exhibited robust antioxidative effects across all extracts, characterized by higher levels of phenolic and flavonoid content. The FT-IR spectrum indicated the presence of compounds such as triglycerides, sterol esters, phospholipids, and unsaturated fatty acids in the extracts.

Conclusion: Collectively, these outcomes prompt the consideration of utilizing the aqueous extract of *Cucurbita maxima* to strengthen humoral and cellular immunity, as well as lymphocyte toxicity, in various ailments.

Keywords: *Cucurbita maxima*, Cell proliferation, IgG production, Immunostimulation, Phagocytosis, Splenocytes, Antioxidant, FT-IR spectrum.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Faculty of Sciences & Techniques, Laboratory of Microbial Biotechnology & Bioactive Molecules, Sidi Mohammed Ben Abdellah University, PO Box 2202, Road of Immouzer, Fez, Morocco; E-mail: aarab.lotfi0@gmail.com

Cite as: Laaraj S, Ouahidi I, Al Moudani N, Allali M, Aarab L. Immunostimulant and Pharmacological Activities, of *Cucurbita Maxima* Seeds on Humoral and Cellular Immunity Cells and their Functions. Open Public Health J, 2024; 17: e18749445274775. <http://dx.doi.org/10.2174/0118749445274775231204095205>



CrossMark

Received: August 29, 2023
Revised: November 01, 2023
Accepted: November 08, 2023
Published: January 31, 2024



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

Immunomodulation represents a therapeutic strategy directed at the targeted adjustment of innate or adaptive immune system components using pharmacological agents or other substances, aiming to achieve either positive

(immunostimulatory) or negative (immunosuppressive) effects. Consequently, compounds of either natural or synthetic origin that alter the immune response, whether generally or specifically through stimulatory or suppressive actions, are categorized as immunomodulatory drugs [1].

Moroccan culture is replete with examples of the utilization of pharmacologically active natural products [2], many of which were employed as immunomodulators. Furthermore, the emergence of oxidative stress triggered by free radicals or reactive oxygen species has been linked to severe health implications. This underscores the importance of investigating the biological impacts of medicinal plant extracts to unearth novel, safe, and potent bioactive compounds possessing immunomodulatory, antibacterial, and antioxidant properties.

Cucurbita Maxima, commonly known as pumpkin, has been recognized as a functional food, ranking among the most popular vegetables globally. Its seeds are abundant sources of bioactive agents boasting nutraceutical attributes. Pumpkin seeds are characterized by high contents of protein, fiber, and fat, reaching percentages of 34.56%, 2.91%, and 36.70%, respectively, alongside relatively lower proportions of sugars and starch, accounting only for 1.08% and 2.15%, respectively [3-5]. Notably, these seeds have numerous biologically active substances that exert substantial pharmacological effects, including antioxidant [6], anti-inflammatory, analgesic [7], and hepatoprotective properties [8].

While prior investigations have delved into the chemical composition, antioxidant capacity, and antibacterial properties of different pumpkin varieties worldwide [3-6], studies focused on the biological characteristics of Moroccan cultivars remain scarce. With this in mind, the present study was conceived to explore not only the immunomodulatory influence of *Cucurbita* seeds on a cell culture model of immune cells through the evaluation of immune function (such as antibody response, T-lymphocyte toxicity, and the complement system) but also to scrutinize the *in vitro* antioxidant potential, look into the functionality of active components via FT-IR spectroscopy, and investigate the antibacterial aspect. These efforts are geared towards uncovering potential applications in the realms of both food and pharmaceutical industries.

2. MATERIALS AND METHODS

2.1. Preparation of Extracts

Seeds of *C. maxima* (pumpkin) were procured from a local market, washed twice with distilled water, and then desiccated at 40°C until a consistent weight was attained. Subsequently, they were finely ground for the purpose of defatting. The defatted powder underwent extraction *via* maceration, utilizing water, ethanol (EtOH), or ethyl acetate (EtOAc) at a concentration of 10%. After filtration, the water extract underwent a 24-hour process of lyophilization. The ethanol and ethyl acetate extracts were evaporated using a rotary evaporator. Finally, all extracts were preserved at -20°C until their intended use.

2.2. Cell Culture

Cell suspensions employed in this study were derived from rabbits that were humanely euthanized. Rabbits weighing between 1.5 to 2 kg were euthanized under petroleum-ether anesthesia. All experimental procedures

adhered to ethical regulations and followed animal care guidelines stipulated by the local Ethics Committee CEFST and approved under reference number 13 /2021/CEFST.

The immune organs, namely the spleen and thymus, were aseptically excised from the animals. Cell suspensions were prepared by gently pressing the organs through a fine wire mesh, following the methodology outlined in our previous research [9-12]. The cells were subsequently washed with RPMI medium, and red blood cells were lysed using 154 mM ammonium chloride. The viability of cells was assessed through microscopic examination employing a 0.1% trypan blue exclusion test.

The culture medium used was RPMI supplemented with 2 mM glutamine, 10% serum, antibiotics (ampicillin 100 U/mL and streptomycin 100 mg/mL), and antifungal agent (Fluconazole 2 mg/mL).

2.3. Cell Proliferation Assay

Cell proliferation was assessed using the MTT assay following the protocol by Mosmann et al., 1983, as previously outlined [11-13]. In brief, cells were seeded at a density of 150,000 cells per well in 96-well plates and subsequently incubated at 37°C within a humidified chamber under an atmosphere consisting of 95% air and 5% CO₂ for a duration of 72 hours. Prior to the incubation, extracts at the desired concentrations were introduced to the cells and diluted in RPMI medium. Following this, a 10 µL volume of MTT solution (5 mg/mL in PBS) was administered. After three hours of incubation, DMSO was introduced to all wells to dissolve the generated formazan. Finally, the spectrophotometer (BioTek L800) was employed to measure the optical density at a wavelength of 570 nm.

2.4. Isolation of Macrophages and Evaluation of their Proliferation and Activity

To outline, macrophages were isolated from spleen cells based on their capacity to adhere to plate wells. To achieve this, 100 µL of a spleen cell suspension at a concentration of 2.10⁶ cells/mL was added to individual wells of a 96-well plate, followed by an incubation period of 2 hours at 37°C to facilitate macrophage adherence. Subsequently, non-adherent cells were removed, and each well was subjected to two washes with sterile RPMI.

For the assessment of macrophage proliferation, adherent macrophages were incubated with various extracts in RPMI. The evaluation of proliferation was conducted using the MTT assay, as previously described.

The phagocytosis assay was conducted according to our prior studies [9, 11-13], employing neutral red. Specifically, adherent macrophages within each well were incubated with 100 µL of RPMI supplemented with 0.075% neutral red. Subsequently, 10 µL of plant extracts were introduced, and the plates were incubated for 2 hours. Ultimately, after discarding the supernatant and rinsing the cells, the reaction was terminated by employing a solution composed of acetic acid (1M) and ethanol (1:1

v/v), which effectively dissolved the phagocytosed neutral red. The degree of phagocytic activity was quantified by measuring the absorbance at 540 nm, corresponding to the peak absorption of the utilized neutral red in the assay.

2.5. Allogenic Mixed Lymphocyte Reaction (MLR)

Freshly isolated thymocytes obtained from the thymus were subjected to incubation with Chicken Red Blood Cells (CRBC) at a ratio of 10^5 CRBC to 10^6 thymocytes in RPMI medium supplemented with serum. This mixture was then incubated for 24 hours at 37°C in a humidified environment, with the extracts either present or absent. The cytotoxicity of thymocytes against CRBC was evaluated by detecting the lysis of CRBC in the medium through measurement of absorbance at 540 nm.

2.6. Complement Test

The complement test was conducted through the assessment of lysis of mouse red blood cells (MRBC) by means of the complement pathway, aided by the presence of antibodies against MRBC. These antibodies were generated by immunizing rabbits using Freund's adjuvant and utilizing mouse RBC as antigens. MRBC were incubated in RPMI medium supplemented with the serum that contained specific anti-MRBC antibodies for a duration of 4 hours, while extracts at a concentration of 2 mg/ml were present, and the incubation was maintained at 37°C. Following the incubation period, the samples were subjected to centrifugation, and the absorbance of the supernatants was measured at 540 nm.

2.7. Evaluation of Total IgG Production by ELISA Assay

The ELISA assay was conducted following the previously established procedures [12, 14-16]. Isolated splenocytes were cultured in RPMI with or without extracts for a duration of 72 hours at 37°C. Subsequently, 100 µL of cell culture supernatant was dispensed onto a microtiter plate for the purpose of IgG assessment. Subsequent to this, a peroxidase-labeled anti-IgG rabbit antibody was introduced (100 µL per well) and incubated for two hours at 37°C. The immune complex was revealed through the addition of the chromogen orthophenylenediamine (OPD), at a concentration of 0.5 mg/mL. The reaction was halted using 3 M HCl, and the absorbance was measured at 490 nm.

For the specific IgG assay, the splenocyte culture was initiated by adding Ovalbumin to RPMI, both with and without extracts. The assay commenced by coating the wells with 100 µL of Ovalbumin (1 g/mL). Following this, 100 µL of the cell culture supernatant was introduced, and the assay was executed in the same manner as outlined for Total IgG.

2.8. Antimicrobial Activity (Agar Well Diffusion Method)

Cucurbita seed extracts were evaluated against a

panel of four Gram-positive bacteria, *Bacillus subtilis* ILP1428B and *Staphylococcus aureus* CIP543154 (Pasteur Institute Collection), as well as five Gram-negative bacteria: *Pseudomonas aeruginosa* ATCC27653 and four strains of *Escherichia coli* CIP5412 (from the American Type Culture Collection).

The surface of agar plates was inoculated by evenly spreading 100 µl of the microbial inoculum across the entire agar surface. Subsequently, an aseptic perforation of 6 to 8 mm diameter was made using a sterile probe or nozzle, and 50 µl of the samples were introduced into the well. The agar plates were then incubated at 37°C for an additional 24 hours. The diameter of inhibition zones was calculated as previously described [17, 18].

2.9. Total Phenolic Content

The total phenolic content of the extract was quantified using the Folin-Ciocalteu method. This technique involved the blending of 1.5 ml of Folin-Ciocalteu reagent (10%) with 200 µl of extracts previously dissolved in methanol. The resulting mixture was gently stirred and allowed to react for 5 minutes in darkness, followed by the addition of 1.5 ml of sodium carbonate solution (5%). Following a 2-hour incubation period at room temperature and in the absence of light, the measurements were taken at 750 nm. The concentration of phenolic content was determined based on a calibration curve utilizing Gallic acid. The outcomes were presented as milligrams equivalent of Gallic acid per gram of extract.

2.10. Total Flavonoid Content

The quantification of flavonoids was conducted using the Aluminum trichloride (AlCl₃) method as described by Bahorun et al., 1996 [19-35]. In this procedure, precisely 0.5 ml of each extract (dissolved in methanol) was combined with 0.1 ml of aluminum chloride (10%), 0.1 ml of potassium acetate (1 M), and 4.3 ml of distilled water. Following shaking and an incubation period of 30 minutes at room temperature, absorbance readings were taken at 415 nm. The concentration of flavonoids was determined through reference to a calibration curve established using quercetin. The results were expressed as micrograms of quercetin equivalent per milligram of extract.

2.11. Total Antioxidant Activity

The DPPH radical scavenging assay was performed following the previously established procedure with slight modifications [10, 19, 20]. In essence, the extracts were diluted in methanol and subsequently combined with a DPPH solution. Absorbance readings were taken at 517 nm.

The FRAP assay was executed according to the method outlined by Oyaizu [21]. Each extract was mixed with 2.5 ml of a 0.2 M phosphate buffer solution (pH 6.6) and 2.5 ml of a 1% potassium ferricyanide solution (K₃Fe(CN)₆). Subsequently, trichloroacetic acid at a concentration of 10% was introduced to halt the reaction. An aliquot (2.5 ml) of the supernatant, diluted with 2.5 ml of distilled water, was combined with 0.5 ml of FeCl₃ solution, and

the absorbance was measured at 700 nm.

The ABTS assay was carried out as detailed by Re et al. [22]. ABTS was dissolved in distilled water, followed by the addition of potassium persulfate. The resultant mixture was left at room temperature in the dark for a duration of 12 to 16 hours before the introduction of the extracts. Subsequently, 2 ml of the diluted ABTS+ solution was added to 200 μ l of each extract and allowed to stand for 1 minute, and the absorbance was determined at 734 nm.

2.12. UV Absorbance

The UV absorbance of the three extracts was assessed in vitro by creating diluted samples with concentrations of 2 mg/ml in ether. Subsequently, absorbance values were ascertained across the wavelength range from 290 nm to 400 nm, which encompasses the UVB and UVA regions. Under identical conditions, we also determined the UV absorbance of two commercial sunscreens to serve as positive controls.

2.13. Infrared Spectral Analysis

Infrared analysis was conducted using the VERTEX 70 - BRUKER infrared spectrophotometer. A small droplet of the purified extract was cautiously positioned onto the

surface of a crystal and secured within the path of the infrared beam. The obtained infrared data was subsequently compared to a table of IR frequencies for analysis.

2.14. Statistical Analysis

Each experimental condition was replicated a minimum of three times (n=3). The provided values are presented as the means \pm standard error of the mean (SEM). Statistical analyses were performed using the Student's t-test. Significance was determined at a threshold of $p < 0.05$.

3. RESULTS

3.1. Effect of *C. maxima* Extracts on Humoral Immunity (Fig. 1, Tables 1, 2)

Fig. (1) illustrates the impact of *C. maxima* extracts obtained through ethanol (EtOH), ethyl acetate (EtOAc), and water on splenocyte proliferation. It is evident that both EtOH and EtOAc extracts significantly inhibited splenocyte proliferation (with inhibitions of 43.2% and 40.5%, respectively, at a concentration of 2 mg/ml). In contrast, the aqueous extract did not alter this proliferation.




Fig. (1). Effect of *C. Maxima* seeds extracts on splenocyte proliferation. EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.

Table 1. Effect of *C. maxima* extracts (at 2 mg/ml) on total igt production results represented in % of control obtained in same experiments.

Cucurbita Extracts		
EtOH(N=6)	EtOAc (N=6)	Water (N=6)
50.5 ± 5.0%	98.9 ± 10.6%	100.1 ± 6.7%

Abbreviations: EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.

Furthermore, we proceeded to assess the total IgG production by splenocytes under the influence of distinct *C. maxima* extracts. As depicted in Table 1, the outcomes indicate that only the EtOH extract of *C. maxima* demonstrated inhibition of Total IgG production (Table 1: exhibiting a 49.5% inhibition).

Lastly, the study delved into examining the complement activity under these various extracts. Notably, the aqueous extract exhibited a slight, though statistically non-significant, stimulation of complement activity, while both the EtOH and EtOAc extracts appeared to induce a minor suppression of complement activity (Table 2).

3.2. Effect of *C. maxima* Extracts on Cellular Immunity (Fig. 2 Table 3)

We conducted an assessment of the extracts’ impact on thymocyte proliferation. As demonstrated in Fig. (2), it is evident that both the EtOH and EtOAc extracts of *C. maxima* elicited inhibition of thymic cell proliferation (with inhibitions of 48.2% and 41.1%, respectively). Conversely,

under aqueous extraction, a significantly pronounced stimulation of thymic cell proliferation was observed, presenting a remarkable 285.7% response (p<0.001; N=5, Fig. 2).

The evaluation of thymic cell cytotoxicity against foreign cells (MLR), as outlined in Table 3, showthat neither the EtOH nor EtOAc extracts induced significant changes in MLR. However, the aqueous extract demonstrated a notably heightened stimulation of cytotoxicity activity (146.5% response compared to control; p<0.005; Table 3).

3.3. Effect of *C. maxima* on Macrophage Proliferation and their Activity (Fig. 3 and Table 4)

In terms of macrophage proliferation, we noted that all *C. maxima* extracts induced marginal stimulation of macrophage proliferation. The highest response was achieved with the EtOH extract, resulting in a maximum response of 187.4% (Fig. 3).




Fig. (2). Effect of *C. Maxima* seeds extracts on Thymocyte proliferation. EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.

Table 2. Effect of *C. maxima* extracts (at 2 mg/ml) on complement activity.

Cucurbita Extracts		
EtOH(N=6)	EtOAc (N=6)	Water (N=6)
87.9 ± 9.6	88.7 ± 9.3	119.9 ± 69.2

Abbreviations: EtOH: ethanol extract; EtOAc: ethyl acetate extract; water: aqueous extract.

Table 3. Effect of *C. Maxima* (at 2 mg/ml) on MLR activity results represented in % of control obtained in same experiments.

Cucurbita Extract		
EtOH(N=7)	EtOAc (N=7)	Water(N=7)
99.9 ± 17.7	110.3 ± 11.6	146.5 ± 14.7

Abbreviations: EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.




Fig. (3). Effect of *C. Maxima* seeds extracts on macrophage proliferation.

EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.

Table 4. Effect of *C. Maxima* extracts on macrophage phagocytosis.

N=4	Control	EtOH	EtOAc	Water
Cucurbita	0.25 ± 0.01	0.26 ± 0.04	0.15 ± 0.03	0.11 ± 0.01

Abbreviations: EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.

The assessment of phagocytic activity (Table 4) show that the EtOH extract did not bring about any alteration in phagocytosis. In contrast, both the EtOAc and aqueous extracts demonstrated inhibitory effects on phagocytosis (with inhibitions of 40% and 56%, respectively; $p < 0.05$;

N=4; Table 4).

3.4. Antimicrobial Activity

The findings pertaining to antimicrobial activity have been consolidated in Table 5. The effectiveness of

antimicrobial activity displayed variations based on the specific extract and the targeted microorganism. The aqueous extract emerged as the most potent, exhibiting inhibition potential against *Staphylococcus aureus*, *Pseudomonas aeruginosa* ATCC53 (Gram-negative), and *E. coli* growth. This confirms the immunostimulatory impact of this extract and its potential utilization as a functional food.

3.5. Total Phenolic and Flavonoid Content

Table 6 presents a summary of the outcomes regarding the total phenolic content derived from *C. maxima* extracts. The observations highlight the higher levels of phenolic and flavonoid content for EtOH and EtOAc extracts compared to the aqueous extracts.

3.6. The Antioxidative Effect of *C. Maxima* Extracts

To assess the antioxidant activity of *C. maxima* extracts, three distinct methods were employed: DPPH, FRAP, and ABTS assays.

Fig. (4) illustrates the antioxidative impact of *C.*

maxima extracts as determined by the DPPH assay. Notably, all three extracts demonstrated substantial antioxidant effects, which intensified with increasing concentrations ranging from 0.5 to 50 mg/ml. Particularly, the EtOH and water extracts displayed maximal effects exceeding 60%. The IC_{50} values for EtOH, EtOAc, and water extracts were recorded at 2.5%, 0.52%, and 0.53% respectively. Comparatively, the standard compounds exhibited IC_{50} values of 0.55%, 0.065%, and 0.085% for Vitamin E, BHT, and ascorbic acid.

Furthermore, the antioxidant power of *C. maxima* extracts was evaluated through the FRAP assay. As indicated in Fig. (5), the EtOAc extract showcased lower reducing power, while the EtOH and aqueous extracts exhibited robust antioxidant potential at concentrations of 5 and 50 mg/ml.

The antioxidant ABTS assay revealed the potent ability of all the analyzed extracts to effectively quench ABTS+. Notably, the aqueous extract stood out with the highest activity, displaying an IC_{50} of 7.5 μ g/ml (Table 7).




Fig. 4 contd.....




Fig. (4). Antioxidative effects of *C.maxima* extracts By DPPH radical scavenging essay compared to BHT, Ascorbic acide and vit E.

Table 5. Antimicrobial activity of *C.maxima* extracts evaluated by agar well diffusion method.

Bacteria	<i>C. maxima</i> Extracts		
	EtOH	EtOAc	Aqueous
Staphylococcus Aureus Cip (gram+)	+	+	10
Bacillus subtilis ILP14 (gram+)	-	-	-
Bacillus subtilis ILB10 (gram+)	+	-	8
Pseudomonas aeroginosae ATCC53 (gram-)	12-11	-	10
E.coli DH5α (gram-)	-	-	-
E.coli cip 5412 (gram-)	+	+	10
E.coli AL52 (gram-)	-	-	-
E.coli HB101 (gram-)	-	-	-
E.coli B12(gram-)	+	-	8

Note: Results are expressed as diameters of growth inhibition zones (mm) including the hole diameter (6.00 mm); (--) No effect.

Table 6. Total phenolic, flavonoid. The values are the mean of three determinations ± standard error.

Samples	Total phenolic content (mg of GA/g Extract)	Total flavonoid content (mg of Q/g Extract)
EtOH	686.18± 0.090	148.6± 0.070
EtOAc	790.54±0.0190	183.4±0.090
Water	627.62±0.100	92.9±0.041

Abbreviations: EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.

Table 7. Antioxidant activity of extracts as determinate by ABTS assay. The values are the mean of three determinations \pm standard error.

Samples	ABTS IC50 (mgTE/edw)
EtOH	9.97 \pm 0.250
EtOAc	11.45 \pm 0.277
Water	7.50 \pm 0.082

Abbreviations: EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.




Fig. (5). The antioxidative effect of *C.maxima* extracts by FRAP reducing power essay.

EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.

3.7. UV Absorbance

Fig. (6) depicts the peak UV absorbance of *C. maxima* extracts in comparison to sunscreens SPF80 and SPF30. In the UVB range, it was noted that the EtOH extract exhibited over 1.5 times higher absorption than that of sunscreen SPF30. This was followed by the aqueous extract with 1.5, while the EtOAc extract exhibited the lowest level of UVB absorption, reaching a maximum of 1.

3.8. Infrared Spectral Analysis

The FTIR spectrum was utilized to identify the functional groups of active components, discerning peak values within the infrared radiation region. The FT-IR spectra of aqueous, ethanolic, and ethyl acetate extracts of *C. maxima* seeds are presented in Figs. (7 to 9),

respectively.

The FT-IR analysis of the aqueous extract unveiled the presence of the aromatic C=C bond observed between 1650 and 1500 cm^{-1} , C—O stretching between 1150 and 1050 cm^{-1} , an out-of-plane O-H bending at 670-640 cm^{-1} , and a robust absorption region between 3500-3000 cm^{-1} attributable to hydroxyl groups, phenolic/acidic groups, or sugar constituents. These functions boost the extract's polar nature. Peaks between 3000 and 2800 cm^{-1} correspond to C-H symmetrical stretching. Additionally, an asymmetric CH_2/CH_3 scissoring at 1420-1470 cm^{-1} and - CH_2 wagging at 880-800 cm^{-1} were identified. These above-mentioned bands potentially demonstrate the presence of triglycerides, sterol esters, or phospholipids in the extracts [23-25].

Furthermore, distinctive bands situated around the C=O group, were evident in the organic extracts. 1750-1740 cm^{-1} (Figs. 8, 9), signifying the stretching of

UV Absorbance




Fig. (6). Maxima absorbance's of *C.maxima* seeds extracts compared to sunscreen of SPF 80 and 30. EtOH: Ethanol extract; EtOAc: Ethyl Acetate extract; Water: Aqueous extract.




Fig. (7). FT-IR analysis of aqueous extract of *C. maxima* seeds.




Fig. (8). FT-IR analysis of ethanolic extract of *C. maxima* seeds.




Fig. (9). FT-IR analysis of ethyl acetate extract of *C. maxima* seeds.

Table 8. IR bands assignments of *C. maxima* seeds extracts with water, EtOH and AcOEt [27, 33].

Frequency. cm ⁻¹	Functional Group v: vibration	Vibration Mode
3650-3200	v(-O-H) (<i>strong</i>) of acidic group, phenolic group and carbohydrate,	stretching
3200-2850	v(C-H)	stretching
1650-1750	v(C-H)	stretching
2950-2800	v(C-H)CH ₃ + v(C-H) CH ₂	stretching
1750	v(C=O) of carbonyl group of ester (<i>strong and broad</i>)	stretching
1650	v(C=O) of acidic / amidic group	stretching
1420-1470	v(C-H) CH ₂ /CH ₃	bending (scissoring)
1350-1250	v(=C-H)	bending
1150	v(C-O)	stretching
1100-1000	v(C-H) CH ₂ /CH ₃	bending
1000-900	v(H-C=C-H)	bending
900-450	v(C-H) CH ₂	bending, rocking, wagging
670-640	v(O-H)	bending

However, the FT-IR analysis of the EtOAc and EtOH extracts yielded analogous results, both displaying a weak polar nature. Upon spectral examination, an absorption band emerged at 1750 cm⁻¹, indicative of a robust C=O absorption band attributable to the carbonyl ester group [23]. Additionally, the intense peaks at 2850-2860 and 2920-2930 cm⁻¹ corresponded to the symmetric and asymmetric stretches of the CH₂ group, while the peak within 2980-3050 cm⁻¹ represented the scissoring vibrations of the CH₂ group. All of these characteristics are attributed to unsaturated fatty acids, triglycerides, or unsaturated lipids present in the extracts. This phenomenon was more pronounced in the case of non-polar character extracts such as EtOAc and EtOH than with the aqueous extract [23-25]. The length of the hydrocarbon chains exceeded or were equal to 4 carbon atoms, as evidenced by the presence of a rocking CH₂ at 740 cm⁻¹ (Table 8) [26].

4. DISCUSSION

The modulation of immune functions using herbal medicines presents an alternative or complementary approach to conventional therapies for treating infections, immunological disorders, and cancer. Numerous medicinal plants have been evaluated for their immunomodulatory activity in humans and animals, both *in vivo* and *in vitro*. The results have consistently demonstrated their beneficial impact on immune system modulation through various mechanisms.

In the Moroccan context, seeds of *C. maxima* have been utilized in folk medicine to address various ailments. Consequently, the aim of our study was to assess the immunomodulatory effects of three extracts of *C. maxima* seeds on immune functions.

The EtOH extract of *C. maxima* inhibited Total IgG production, whereas the aqueous extract remarkably stimulated complement activity, suggesting the potential of *C. maxima* to modulate humoral immune responses. These findings align with previous research by [27], where

C. maxima seed-treated rabbits exhibited increased antibody titers, total white blood cell counts, and serum immunoglobulins in both normal and suppressed animals. Furthermore, *C. maxima* seeds are renowned for their abundance in bioactive compounds, such as carotenoids, tocopherols, and sterols, which are useful for a broad spectrum of biological activities, as confirmed by *in vivo* experiments [28, 29].

For cellular immunity, aqueous extracts of *C. maxima* seeds demonstrated higher stimulation of thymic cell proliferation and Major Histocompatibility Complex (MHC) class II Restricted T-cell proliferation (MLR) activity. This trend is consistent with findings by [27], affirming the potential of *C. maxima* seeds against cell-mediated immunosuppression in rabbits.

Our observations also revealed that EtOAc and aqueous extracts of *C. maxima* seeds induced a reduction in the phagocytic function of macrophages. This outcome can be attributed to the anti-inflammatory activity of these extracts, in line with the results obtained by [7]. When evaluating the safety profile and exploring the analgesic and anti-inflammatory activity of ethanol extract of *Cucurbita maxima* seeds, the seed extract of *C. maxima* was found to be safe and showed significant analgesic and anti-inflammatory activity in comparison with the control group. The results were similar to the study of [30]. *Cucurbita moschata* Dush, a plant belonging to the Cucurbitaceae family, was examined in mouse splenocytes and macrophage cell lines. The primary component in Dush is β-carotene. According to the results, both have an immune-enhancing effect by activating macrophages and splenocytes to produce Th1 cytokines. This suggests that β-carotene has an anti-inflammatory effect by significantly upregulating the production of TNFα, IFNγ, and IL2 Choi et al. [31]. This demonstrates the implication of all metabolites in anti-inflammation pathways. In contrast [36], reports that polyose extract from pumpkin can increase the weight of the immune function of mice as well as the phagocytic function of the macrophages.

Additionally, a preliminary antimicrobial screening of

C. maxima seed extracts demonstrated inhibitory effects against *Staphylococcus aureus* (gram+), *Pseudomonas aeruginosa* ATCC53 (gram-), and *E. coli* B12 (gram-), in line with earlier reports on pumpkin (*Cucurbita pepo* L.) seed oil [32].

The total phenolic content ranged from 627.62 ± 0.1 to 790.54 ± 0.019 mg of gallic acid per gram of extract. The order of total phenolic content was EtOAc > EtOH > Water (Table 5), echoing the findings of [6], who examined a variety of pumpkin seeds and identified high content and diversity of polyphenols. In another study by Pazinato et al. [33], the phenolic compounds content was shown to be between 611.39 and 1053.18 (mg GAE per 100 g of oil). Furthermore, the flavonoid content in our study was lower than that reported by [31] for *C. maxima* juices.

The antioxidative effect results were congruent with the research of Nawirska-Olszańska et al. [6], who examined the antioxidative effect of the seeds of 12 pumpkin varieties belonging to the species *Cucurbita maxima* Duch. and *Cucurbita pepo* L. The seeds of *C. pepo* exhibit better antioxidant properties regardless of the extraction solvent used. The same result was found for other studies [33], for which the IC50 values were 554.2 to 788.8 (μg per ml). For the oil extracted from the mixture (seeds/peel), while our results were superior to those obtained by Rezig et al. [3], 36.22% of DPPH radicals were quenched by pumpkin seed oil. In addition, two varieties of pumpkin seed oils were potentially active. IC50 of indigenous pumpkin seed oil (IPSO) and hybrid pumpkin seed oil (HPSO) were observed as 3.72 ± 0.93 $\mu\text{g}/\text{mL}$ and 5.73 ± 0.96 $\mu\text{g}/\text{mL}$, respectively [30].

In summary, our findings align with previous studies, providing further the remarkable antioxidant properties of *C. maxima* seeds and the potential for photoprotection offered by the EtOAc extract which is in line with our previous study [10]. This photoprotection aspect deserves further investigation. Currently, this article uses *in vitro* tests, and future research will include *in vivo* tests to supplement it.

CONCLUSION

In conclusion, our research findings underscore the immune-enhancing potential of aqueous *C. maxima* extract, manifesting through the stimulation of spleen cell proliferation, complement activity, thymic cell function, and cytotoxicity. These results suggest the potential utility of aqueous *Cucurbita* extract for boosting both humoral and cellular immunity, along with lymphocyte toxicity modulation across various diseases. Furthermore, these extracts exhibit substantial antioxidant and antimicrobial activities, in line with their high total phenolic and flavonoid content. Additionally, their capacity for UV ray absorption enhances their potential as photoprotective agents.

LIST OF ABBREVIATIONS

CRBC = Chicken Red Blood Cells

FT-IR = Fourier-transform Infrared Spectroscopy

ETHICAL STATEMENT

All experimental procedures adhered to ethical regulations and followed animal care guidelines stipulated by the local Ethics Committee CEFST. under reference no. 13/2021/CEF51 morocco.

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

The data and supportive information are available within the article.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Mukherjee S, Mukherjee U. A comprehensive review of immunosuppression used for liver transplantation. *J Transplant* 2009; 2009: 701464. <http://dx.doi.org/10.1155/2009/701464>
- [2] Bellakhdar J, Claisse R, Fleurentin J, Younos C. Repertory of standard herbal drugs in the Moroccan pharmacopoea. *J Ethnopharmacol* 1991; 35(2): 123-43. [http://dx.doi.org/10.1016/0378-8741\(91\)90064-K](http://dx.doi.org/10.1016/0378-8741(91)90064-K) PMID: 1809818
- [3] Rezig L, Chouaibi M, Msaada K, Hamdi S. Chemical composition and profile characterisation of pumpkin (*Cucurbita maxima*) seed oil. *Ind Crops Prod* 2012; 37(1): 82-7. <http://dx.doi.org/10.1016/j.indcrop.2011.12.004>
- [4] Stevenson DG, Eller FJ, Wang L, Jane JL, Wang T, Inglett GE. Oil and tocopherol content and composition of pumpkin seed oil in 12 cultivars. *J Agric Food Chem* 2007; 55(10): 4005-13. <http://dx.doi.org/10.1021/jf0706979> PMID: 17439238
- [5] Montesano D, Blasi F, Simonetti M, Santini A, Cossignani L. Chemical and nutritional characterization of seed oil from *Cucurbita maxima* L. (Var. Berrettina) pumpkin. *Foods* 2018; 7(3): 30. <http://dx.doi.org/10.3390/foods7030030> PMID: 29494522
- [6] Nawirska-Olszańska A, Kita A, Biesiada A, Sokół-Łętowska A, Kucharska AZ. Characteristics of antioxidant activity and composition of pumpkin seed oils in 12 cultivars. *Food Chem* 2013; 139(1-4): 155-61. <http://dx.doi.org/10.1016/j.foodchem.2013.02.009> PMID: 23561092
- [7] Wahid S, Alqahtani A, Alam Khan R. Analgesic and anti-inflammatory effects and safety profile of *Cucurbita maxima* and *Cucumis sativus* seeds. *Saudi J Biol Sci* 2021; 28(8): 4334-41. <http://dx.doi.org/10.1016/j.sjbs.2021.04.020>
- [8] Abou Seif HS, Seif A. Ameliorative effect of pumpkin oil (*Cucurbita pepo* L.) against alcohol-induced hepatotoxicity and oxidative stress in albino rats. *Beni Suef Univ J Basic Appl Sci* 2014; 3(3): 178-85. <http://dx.doi.org/10.1016/j.bjbas.2014.08.001>
- [9] Laaraj S, Ouahidi I, Aarab L. Immunostimulant effect of *Brassica rapa* and *Raphanus sativus* seeds on thymic cells and their cytotoxicity. *Egypt J Immunol* 2023; 30(1): 87-95. <http://dx.doi.org/10.55133/eji.300109> PMID: 36592121
- [10] Laaraj S, Ouahidi IAL. Photoabsorbent, antioxidative, and anti-

- inflammatory properties of three traditional cosmetic seed oils. Trop J Nat Prod Res 2022; 6(11): 1790-3.
- [11] Daoudi A, Abdel-Satter E, Aarab L. The relationship between lectin compounds and immunomodulatory activities of protein extracted from plants. J Plant Stud 2013; 3(1): 56-64. <http://dx.doi.org/10.5539/jps.v3n1p56>
- [12] El Hamsas El Youbi A, Bousta D, Ouahidi I, Aarab L. Criblage pharmacologique primaire d'une plante endémique originaire du Sud Marocain (*Tetraena gaetula* [Emb. & Maire] Beier & Thulin). C R Biol 2010; 333(10): 736-43. <http://dx.doi.org/10.1016/j.crvi.2010.08.001> PMID: 20965443
- [13] Daoudi A, Bagrel D, Aarab L. *In vitro* anticancer activity of some plants used in Moroccan traditional medicine. J Med Plants Res 2013; 7(17): 1182-9.
- [14] Amal B, Najlae M, Ouarda A. Immunomodulating properties of protein fractions isolated from Delphinium staphysagria seeds. J Med Plants Res 2016; 10(3): 29-34. <http://dx.doi.org/10.5897/JMPR2015.5957>
- [15] Mejrhit N, Azdad Y, Azdad O, Aarab L. Determination of histamine levels in commonly consumed fish in the region of Fez. Br Food J 2018; 120(10): 2388-94. <http://dx.doi.org/10.1108/BFJ-12-2017-0670>
- [16] Ouahidi I, Barakat A, El Youbi El Hamsas A, Aarab L. Variation de l'immunoréactivité des IgE et des IgG vis-à-vis des gliadines sous l'effet du pH et de la température. Immunol Anal Biol Spec 2011; 26(2): 68-75. <http://dx.doi.org/10.1016/j.immbio.2010.10.005>
- [17] Elaaraj I, Er S, Nakkabi A, Es-sounni B, Koraichi I, El N, et al. Journal of the indian chemical society synthesis, characterization and antioxidant, antibacterial activity Zn²⁺. J Indian Chem Soc 2022; 99(5): 100404. <http://dx.doi.org/10.1016/j.jics.2022.100404>
- [18] Balouiri M, Sadiki M, Ibsouda SK. Methods for *in vitro* evaluating antimicrobial activity: A review. J Pharm Anal 2016; 6(2): 71-9. <http://dx.doi.org/10.1016/j.jpha.2015.11.005> PMID: 29403965
- [19] Taouda H, Alaoui MM, Errachidi F, Chabir R, Aarab L. Comparative study of the morpho-metric and Biochemical dates caractere solding in the regional market of FES / MORO 2014; 8(1): 1-10.
- [20] Boudkhili M, Greche H, Misbahi H, et al. Isolation and antioxidant activity of flavonoids from *Coriaria myrtifolia* methanolic extract. Chem Nat Compd 2015; 51(1): 141-2. <http://dx.doi.org/10.1007/s10600-015-1222-y>
- [21] Oyaizu M. Studies on products of browning reaction. Antioxidative activities of products of browning reaction prepared from glucosamine. Eiyogaku Zasshi 1986; 44(6): 307-15. <http://dx.doi.org/10.5264/eiyogakuzashi.44.307>
- [22] Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. Free Radic Biol Med 1999; 26(9-10): 1231-7. [http://dx.doi.org/10.1016/S0891-5849\(98\)00315-3](http://dx.doi.org/10.1016/S0891-5849(98)00315-3) PMID: 10381194
- [23] Bouramdane Y, Fellak S, El Mansouri F. Impact of Natural Degradation on the Aged Lignocellulose Fibers of Moroccan Cedar Softwood : Structural Elucidation by Infrared Spectroscopy (ATR-FTIR) and X-ray Diffraction. XRD 2022.
- [24] Boukir A, Fellak S, Doumenq P. Structural characterization of *Argania spinosa* Moroccan wooden artifacts during natural degradation progress using infrared spectroscopy (ATR-FTIR) and X-Ray diffraction (XRD). Heliyon 2019; 5(9): e02477. <http://dx.doi.org/10.1016/j.heliyon.2019.e02477> PMID: 31687572
- [25] Boukir A, Mehyaoui I, Fellak S, Asia L, Doumenq P. The effect of the natural degradation process on the cellulose structure of Moroccan hardwood fiber: A survey on spectroscopy and structural properties. Mediterranean Journal of Chemistry 2019; 8(3): 179. <http://dx.doi.org/10.13171/mjc8319050801ab>
- [26] Boukir A, Guiliano M, Doumenq P, El A, Mille G. Caractérisation structurale d'asphaltes bitumineux par spectroscopie infrarouge (IRTF). Application à la photo-oxydation 1998; 597-602.
- [27] Ranganathan V, Selvasubramanian S, Vasanthakumar S. Estimation of humoral immune response in rabbits fed with *Cucurbita maxima* seeds. Veterinary World 2013; 6(7): 396. <http://dx.doi.org/10.5455/vetworld.2013.396-399>
- [28] Dyshlyuk L, Babich O, Prosekov A, Ivanova S, Pavsky V, Yang Y. *In vivo* study of medical and biological properties of functional bakery products with the addition of pumpkin flour. Bioactive Carbohydrates and Dietary Fibre 2017; 12(January): 20-4. <http://dx.doi.org/10.1016/j.bcdf.2017.09.001>
- [29] Patel S. Pumpkin (*Cucurbita* sp.) seeds as nutraceutical: a review on status quo and scopes. Med J Nutrition Metab 2013; 6(3): 183-9. <http://dx.doi.org/10.3233/s12349-013-0131-5>
- [30] Amin MZ, Rity TI, Uddin MR, Rahman MM, Uddin MJ. A comparative assessment of anti-inflammatory, anti-oxidant and anti-bacterial activities of hybrid and indigenous varieties of pumpkin (*Cucurbita maxima* Linn.) seed oil. Biocatal Agric Biotechnol 2020; 28(June): 101767. <http://dx.doi.org/10.1016/j.bcab.2020.101767>
- [31] Choi KC, Cho SW, Kook SH. Intestinal anti-inflammatory activity of the seeds of *Raphanus sativus* L. in experimental ulcerative colitis models. J Ethnopharmacol 2016; 179: 55-65. <http://dx.doi.org/10.1016/j.jep.2015.12.045> PMID: 26721217
- [32] Bardaa S, Ben Halima N, Aloui F. Oil from pumpkin (*Cucurbita pepo* L.) seeds: Evaluation of its functional properties on wound healing in rats. Lipids Health Dis 2016; 15(1): 73. <http://dx.doi.org/10.1186/s12944-016-0237-0> PMID: 27068642
- [33] Cuco RP, Cardozo-Filho L, Silva C. Simultaneous extraction of seed oil and active compounds from peel of pumpkin (*Cucurbita maxima*) using pressurized carbon dioxide as solvent. J Supercrit Fluids 2019; 143(143): 8-15. <http://dx.doi.org/10.1016/j.supflu.2018.08.002>
- [34] Abirami A, Nagarani G, Siddhuraju P. *In vitro* antioxidant, anti-diabetic, cholinesterase and tyrosinase inhibitory potential of fresh juice from *Citrus hystrix* and *C. maxima* fruits. Food Sci Hum Wellness 2014; 3(1): 16-25. <http://dx.doi.org/10.1016/j.fshw.2014.02.001>
- [35] Baharun T, Gressier B, Trotin F, et al. Oxygen species scavenging activity of phenolic extracts from hawthorn fresh plant organs and pharmaceutical preparations. Arzneimittelforschung 1996; 46(11): 1086-9. PMID: 8955870
- [36] BAO. Effects of pumpkin polycose and *Capparis spinosa* Linn polycose on immune function of mice. Xinjiang Nongye Kexue 2010; 47(3): 508-11.