



Prevalence and Risk Factors of Hypertension among Young Adults: An Indonesian Basic Health Survey

Laluna Rachma Putri¹ , Mahalul Azam^{1,*} , Alfiana Ainun Nisa¹ , Arulita Ika Fibriana¹ , Phitsanuruk Kanthawee²  and Syed Abdul Shabbir³ 

¹Departement of Public Health, Faculty of Medicine, Universitas Negeri Semarang, Indonesia

²School of Health Science, Mae Fah Luang University, Chiang Rai, Thailand

³Graduate Institute of Biomedical Informatics, Taipei Medical University, Taiwan

Abstract:

Introduction: Hypertension patients aged under 40 years have low awareness and under-diagnosis, hence poor blood pressure control. The prevalence of undiagnosed hypertension in men and women aged 26 to 35 years is 55% and 44%, respectively. This study aimed to provide information on the prevalence and risk factors for hypertension among young adults based on the basic health survey [*Riset Kesehatan Dasar; RISKESDAS*] in Indonesia.

Methods: This cross-sectional study utilized secondary data from the *RISKESDAS* 2018 with a total of 140,073 subjects aged 26-35 years. The characteristics of the subjects were presented as proportion. This study performed a chi-square test and involved multivariate analysis using binary logistic regression for all parameters with p -value < 0.25 in the chi-square test.

Results: This study reports the prevalence of hypertension among young adults to be 4.1%. A study also found that female [POR=1.225 95% confidence interval [CI]: 1.119-1.341], unmarried [POR=0.711; 95% CI:0.556-0.908], low-level education [POR=1.286; 95% CI:1.195-1.384], unemployed [POR=1.267; 95% CI:1.166-1.377], BMI \geq 25 [POR=2.472 [2.295-2.662], diabetes mellitus [POR=3.688; 95% CI:2.630-5.173], emotional mental disorder [POR=2.041; 95% CI:1.856-2.244], instant food consumption [POR=1.108; 95% CI:1.025-1.197] were related to the occurrence of hypertension among young adults.

Conclusion: The prevalence of hypertension among young adults was 4.1%. Females, married, low level of education, unemployed, BMI \geq 25, diabetes mellitus, having a mental disorder, and consumption of instant food altogether increase the risk of getting hypertension among young adults aged 26-35 years old.

Keywords: Hypertension, Risk factors, Young adults, National health survey, Indonesia.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Departement of Public Health, Faculty of Medicine, Universitas Negeri Semarang, Indonesia; E-mail: mahalul.azam@mail.unnes.ac.id

Cite as: Putri L, Azam M, Nisa A, Fibriana A, Kanthawee P, Shabbir S. Prevalence and Risk Factors of Hypertension among Young Adults: An Indonesian Basic Health Survey. *Open Public Health J*, 2024; 17: e18749445361291. <http://dx.doi.org/10.2174/0118749445361291241129094132>



Received: September 30, 2024

Revised: November 03, 2024

Accepted: November 12, 2024

Published: December 10, 2024



Send Orders for Reprints to reprints@benthamscience.net

1. INTRODUCTION

Hypertension is a condition in which repeated examination results in a systolic blood pressure [SBP] of more than 140 mm Hg in the clinic or health center and a

diastolic blood pressure [DBP] of at least 90 mm Hg. These definitions apply to all individuals [>18 years old] [1]. It develops when the arterioles put too much pressure on the vessel walls from the blood, making the heart work harder to keep the pressure [2]. Hypertension can trigger

other non-communicable diseases [3]. The number of hypertension patients continues to increase every year. In 2025, there will be 1.5 billion people with hypertension [4]. Every year, it is estimated that there will be 9.4 million deaths due to hypertension and its complications [2]. Worldwide, there are 54% of adults diagnosed with hypertension [5]. An estimated 46% of adults with hypertension are in ignorance [2]. A rise in blood pressure increases the risk of heart attack, heart failure, stroke, and kidney diseases [6]. Hypertension patients have a 2.6 times greater risk of developing coronary heart disease compared to non-hypertensive individuals [7]. The goal for individuals with hypertension is to maintain BP at <140/90 mm Hg to prevent complications [6].

The prevalence of hypertension in Indonesia has increased, based on the basic health survey [*Riset Kesehatan Dasar; RISKESDAS*]. In 2013, hypertension was 25.8%, while the in 2018 increased to 34.1%. Moreover, there are plenty of unreported cases; merely 1/3 of hypertension cases in Indonesia are diagnosed [8]. Based on a systematic review study in 90 countries, only 46.5% of adults aged ≥ 20 years old with hypertension were aware of their BP [9]. Hypertension patients aged under 40 years have low awareness, under-diagnosis, and poor blood pressure control [10]. The prevalence of undiagnosed hypertension in men and women aged 26 to 35 years is 55% and 44%, respectively. The prevalence of undiagnosed hypertension in young adults is higher than in people aged ≥ 75 years [11].

It is important to identify and manage modifiable risk factors for hypertension. Since the determinants may vary across different age groups, examining the factors specifically among young adults in Indonesia is essential. A limited number of studies have been conducted to investigate the factors contributing to high blood pressure in young adults, especially in Indonesia's 26-35 age group. This study aimed to provide information on the prevalence and risk factors for hypertension among young adults based on the *RISKESDAS* 2018 in Indonesia.

2. MATERIALS AND METHODS

2.1. Design and Study Population

This cross-sectional study utilized secondary data from the *RISKESDAS* 2018. *RISKESDAS* is a population health study conducted by the National Institute for Health Development Policy, Ministry of Health, Republic Indonesia. *RISKESDAS* aims to monitor public health status and services that provide information on key health indicators such as health status, nutritional status, environmental health, health behaviors, and various aspects of health care. *RISKESDAS* data provides health data at district/city and national levels. This report included subjects aged 26-35 who were interviewed by the *RISKESDAS* 2018 enumerator, with a total of 140,073 subjects aged 26-35. Of the 140,073 subjects, 78,300 met the criteria to be analyzed (Fig. 1). Complete detailed reports of the survey methods, including data collection, ethical issues, and other related steps, were published in the *RISKESDAS* 2018 report [12].

2.2. Data Collection and Variables

This study conformed to the ethical criteria based on the Helsinki Declaration of 1975 and its revised in 2013 and approved by the Ethical Review Board, Faculty of Medicine, Universitas Negeri Semarang, under the approval ID number 169/KEPK/FK/KLE/2024. The National Institute for Health Development Policy, Ministry of Health, Republic Indonesia, approved the *RISKESDAS* data for this report. This study used certain parameters from *RISKESDAS* 2018, which were analyzed in this study. These include their weight and height, sociodemographic data, hypertension status, diabetes mellitus status, emotional mental disorder status, smoking status, physical activity status, alcohol consumption, body mass index, salt food consumption, fat food consumption, and instant food consumption.

Sociodemographic factors in this study included age, sex, marital status, residence area, education level, and working status. Hypertension was categorized as previously diagnosed as hypertension by a doctor and having a history of under-hypertension medication. The subjects diagnosed with diabetes mellitus [DM] were classified as having a plasma glucose level of at least 126 mg/dL during fasting or two hours postprandial and a random plasma glucose level of at least 200 mg/dL or a previous medical diagnosis. Moreover, to reduce the possibility of bias concerning the inquiry of "previously had been diagnosed with diabetes mellitus by a physician," inquiries concerning the DM medication and plasma glucose test were asked.

This study used the Ministry of Health of the Republic of Indonesia age category, which young adults defined as someone between 26 and 35 years old [13]. Marital status was divided into three categories: unmarried, married, and widow. The residence areas were divided into two categories: urban and rural. Education level was divided into two categories: low [less than senior high school] and high [more than or equal to senior high school]. Employment status was classified as either employed or unemployed.

BMI was acquired by calculating the result measurement of weight [in kilograms] divided by height [in meters square] and was categorized as ≥ 25 and < 25 . Emotional mental disorders were determined by a self-reporting questionnaire [SRQ] consisting of 20 questions invented by WHO with a cutoff point of 6 [14].

Lifestyle and eating behavior factors in this study included smoking, physical activity, alcohol consumption, salt food consumption, fat food consumption, and instant food consumption. Physical activity was categorized as sedentary and active based on the International Physical Questionnaire. Consumption of salt, fat, and instant food was counted as frequent if consumed ≥ 3 -6 times/week.

2.3. Statistical Analysis

The characteristics of the subjects were presented as proportion. This study performed a chi-square test to find out the risk factors associated with hypertension among young adults in Indonesia. The p -values < 0.05 reflected as statistically significant. This study involved multivariate




Fig. (1). Subjects' recruitment flow.

analysis using a binary logistic regression for all parameters with p -value < 0.25 in the chi-square test. The statistical analysis was accomplished in this study using IBM SPSS Statistics software [for Windows, Version 22.0, Armonk, NY: IBM Corp.].

3. RESULTS

3.1. Characteristics of Subjects

Of the 140,073 subjects aged 26-35, 73,800 met the criteria and were included in the analysis. Among these, 3,225 [4.1%] young adults in Indonesia were identified as having hypertension. Tables 1 and 2 explain the characteristics of these subjects.

Based on the result of the chi-square test in Table 3,

some parameters show p -value < 0.005 ; it refers to evidence of influence hypertension among young adults, *i.e.* sex [POR=0.856; 95% CI: 0.794-0.924], education levels [POR=1.355; 95% CI: 1.262-1.454], marital status [POR=0.612; 95% CI: 0.481-0.777], working status [POR=1.358; 95% CI: 1.264-1.460].

It is also based on the result of the chi-square test in Table 4, body mass index [POR=2.424; 95% CI: 2.255-2.606], diabetes mellitus [POR=4.257; 95% CI: 3.059-5.924], emotional mental disorder [POR=2.105; 95% CI: 1.919-2.311], smoking [POR=0.876; 95% CI: 0.808-0.949], and instant food consumption [POR=1.159; 95% CI: 1.074-1.251]. Binary logistic regression will be used to continue analyzing the variables whose p -value is less than 0.25.

Table 1. The characteristics of subjects.

Variables	Samples [N]	Percentages [%]
Sex	-	-
Male	27227	34.8
Female	51073	65.2
Marital Status	-	-
Unmarried	8398	10.7
Married	67895	86.7
Widowed	2007	2.6
Residence Area	-	-
Urban	34367	43.9
Rural	43933	55.1
Education Level	-	-
Low	37482	33.6
High	51958	66.4

Table 2. The characteristics of subjects.

Variables	Samples [N]	Percentages [%]
Working Status	-	-
Unemployed	26342	33.6
Employed	51958	66.4
Body Mass Index	-	-
BMI \geq 25	31430	40.1
BMI>25	46870	59.9
Diabetes Mellitus	-	-
Yes	274	0.3
No	78026	99.7
Emotional Mental Disorder	-	-
Yes	7705	9.8
No	70595	90.2
Smoking	-	-
Yes	22245	28.4
No	56055	71.6
Physical Activity	-	-
Sedentary	7067	9.0
Active	71233	91.0
Alcohol Consumption	-	-
Yes	3540	4.5
No	74760	95.5
Salt Food Consumption	-	-
Frequent [\geq 3-6 times/week]	32153	41.1
Rarely [$<$ 3-6 times/week]	48147	58.9
Fat Food Consumption	-	-
Frequent [\geq 3-6 times/week]	47346	60.5
Rarely [$<$ 3-6 times/week]	30954	39.5
Instant Food Consumption	-	-
Frequent [\geq 3-6 times/week]	21864	27.9
Rarely [$<$ 3-6 times/week]	56436	72.1
Hypertension	-	-
Yes	3225	4.1
No	75075	95.9

Table 3. The characteristics of subjects based on hypertension status.

Variables	Hypertension				p-value	POR [95% CI]
	Yes		No			
	N	%	N	%		
Sex	-	-	-	-	-	-
Male	1015	3.7	26212	96.3	<0.001*	0.856 [0.794-0.924]
Female	2210	4.3	48863	95.7	-	
Marital Status	-	-	-	-	-	-
Unmarried	253	3.0	8145	97.0	<0.001*	0.612 [0.481-0.777]
Married	2875	4.2	65020	95.8	0.190	0.871 [0.708-1.071]
Widowed	97	4.8	1910	95.2	-	-
Education Level	-	-	-	-	-	-
Low	1778	4.7	35704	95.3	<0.001*	1.355 [1.262-1.454]
High	1447	3.5	39371	96.5	-	
Working Status	-	-	-	-	-	-
Unemployed	1305	5.0	25037	95.0	<0.001*	1.358 [1.264-1.460]
Employed	1920	3.7	50038	96.3	-	

Note: * = p-value <0.05; POR: prevalence odds ratio.

Table 4. The characteristics of subjects based on hypertension status.

Variables	Hypertension				p-value	POR [95% CI]
	Yes		No			
	N	%	N	%		
Body Mass Index	-	-	-	-	-	-
BMI \geq 25	1968	6.3	29462	93.7	<0.001*	2.424 [2.255-2.606]
BMI<25	1257	2.7	45613	97.3	-	
Diabetes Mellitus	-	-	-	-	-	-
Yes	42	15.3	232	84.7	<0.001*	4.257 [3.059-5.924]
No	3183	4.1	74843	95.9	-	
Emotional Mental Disorder	-	-	-	-	-	-
Yes	583	7.6	7122	92.4	<0.001*	2.105 [1.919-2.311]
No	2642	3.7	67953	96.3	-	
Smoking	-	-	-	-	-	-
Yes	835	3.8	21410	96.2	0.001*	0.876 [0.808-0.949]
No	2390	4.3	53665	95.7	-	
Physical Activity	-	-	-	-	-	-
Sedentary	298	4.2	6769	95.8	0.663	1.027 [0.910-1.160]
Active	2927	4.1	68306	95.9	-	
Alcohol Consumption	-	-	-	-	-	-
Yes	159	4.5	3381	95.5	0.260	1.100 [0.934-1.294]
No	3066	4.1	71694	95.9	-	
Salt Food Consumption	-	-	-	-	-	-
Frequent [\geq 3-6 times/week]	1309	4.1	30844	95.9	0.583	0.980 [0.912-1.053]
Rarely [$<$ 3-6 times/week]	1916	4.2	44231	95.8	-	
Fat Food Consumption	-	-	-	-	-	-
Frequent [\geq 3-6 times/week]	1954	4.1	45392	95.9	0.897	1.005 [0.935-1.081]
Rarely [$<$ 3-6 times/week]	1271	4.1	29683	95.9	-	
Instant Food Consumption	-	-	-	-	-	-
Frequent [\geq 3-6 times/week]	995	4.6	20869	95.4	<0.001*	1.159 [1.074-1.251]
Rarely [$<$ 3-6 times/week]	2230	4.0	54206	96.0	-	

Note: * = p-value <0.05; POR: prevalence odds ratio.

Table 5. Binary logistic regression of hypertension risk factors among young adults subject.

Variables	B	S.E	Wald	p-value	POR [95% CI]
Female	0.203	0.046	19.368	<0.001*	1.225 [1.119-1.341]
Unmarried	-0.341	0.125	7.434	0.006*	0.711 [0.556-0.908]
Low Level of Education	0.251	0.037	45.177	<0.001*	1.286 [1.195-1.384]
Unemployed	0.237	0.043	30.930	<0.001*	1.267 [1.166-1.377]
BMI \geq 25	0.905	0.038	573.050	<0.001*	2.472 [2.295-2.662]
Diabetes Mellitus	1.305	0.173	57.182	<0.001*	3.688 [2.630-5.173]
Emotional Mental Disorder	0.713	0.048	216.699	<0.001*	2.041 [1.856-2.244]
Instant Food Consumption	0.102	0.040	6.722	0.010*	1.108 [1.025-1.197]
Constant	-3.844	0.113	1165.952	<0.001	0.021

Note: * = p-value <0.05; POR=prevalence odds ratio.

Table 5 shows the final regression model results, *i.e.*, female [POR:=1.225 95% confidence interval [CI]: 1.119-1.341], unmarried [POR=0.711; 95% CI:0.556-0.908], low-level education [POR=1.286; 95% CI:1.195-1.384], unemployed [POR=1.267; 95% CI:1.166-1.377], BMI \geq 25 [POR=2.472 [2.295-2.662], diabetes mellitus [POR=3.688; 95% CI:2.630-5.173], emotional mental disorder [POR=2.041; 95% CI:1.856-2.244], instant food consumption [POR=1.108; 95% CI:1.025-1.197] were altogether related to the occurrence of hypertension among young adults.

4. DISCUSSION

This national scope study in Indonesia found that the prevalence of hypertension among young adults aged 26-35 was 4.1%. Previous studies reported different prevalences of hypertension among young adults; the CARDIA study reported that in the US, young adults was 9.6% [15]; the Framingham study discovered that the prevalence of hypertension was 12.9% in women and 14.2% in men aged 20 to 49 [16]; another study in the US reported the prevalence of hypertension among 20 to 44 years old was 9.3% [17]. Other studies consistently reported similar prevalence: Albania, aged 15-49, was 16.24% [18]; Zimbabwe, aged 18-25, was 7.4%; Lahore, Pakistan, aged 18-39, was 15.7% [19]; and India, aged 18-39, was 13.8% [20]. This study's findings indicate that the prevalence of hypertension in young adults was low compared to other studies. This is likely since the study only included participants who had received a medical diagnosis or had taken antihypertensive medication. In contrast, previous studies confirmed the diagnosis by blood pressure measurement; missing out on subjects unaware of hypertension was potentially a lot in this study. Most young adults are unaware of their abnormally high blood pressure and only realize its effects when they age [21]. Young adults are less sensitive to hypertension than older people because young adults visit doctors less often than older people [21]. Hypertension patients aged below 40 years have low awareness, under-diagnosis, and poor blood pressure control [10].

This study found that females were more susceptible to hypertension compared to males. Previous studies reported different conditions [18, 22-24], *i.e.*, a study in Albania reported that males are more at risk of

hypertension [18], and a study in Poland also showed similar results [22]. A cross-sectional study conducted in Varanasi revealed that males are at 1.97 times greater risk of hypertension [23]. A study of young adults in Zimbabwe also found that males were 1.53 times more likely to develop hypertension than females [25]. Generally, males have less favorable habits compared to females, such as excess consumption of instant food, smoking, and alcohol consumption, leading to hypertension [18, 26]. However, this study found that females have more risk. The low number of male visits to health services could be the reason for the lower rate of hypertension in males because of undetected cases. Females are more likely to visit health services for consultation regarding birth control and regular gynecological services, where every visit will check their blood pressure so that females' blood pressure is more detected compared to males [21].

This study also found that married persons have a greater risk of hypertension, and unmarried persons have a lower risk. This finding is dissimilar from previous studies that found unmarried persons have a greater risk of hypertension [23, 26, 27]. Another study reported that early-married women have a higher risk of developing hypertension [28]. The present study also found that low-level education subjects possess a higher risk of developing hypertension. A Chinese study found that people with only an elementary education or less are more likely to develop hypertension than people with a middle or higher education [29]. Low knowledge which is a result of low education, influences an unhealthy lifestyle, and access to health information is more accessible for those with higher education levels. The degree of education affects how well they receive and absorb information, which affects their health [23, 30]. Other studies also show that a higher level of education is expected to increase good knowledge about hypertension and its treatment [31].

This study also found that unemployed subjects have a greater risk of hypertension. A previous study concluded similarly that unemployed subjects risk developing hypertensive disease more than employed people [32-34]. A systematic review and meta-analysis study in Southeast Asian Countries also stated that the unemployed have a greater risk of hypertension [34]. Unemployed people are

1.39 more at risk of hypertension compared to employed people [35]. The results of a nationwide study in China state that unemployment conditions are a high risk of hypertension because of the relationship with economic capacity. Therefore, unemployment will indirectly make someone less aware of hypertension treatment and control [36, 37]. Unemployment is more at risk of hypertension because of low income; a person will have a limited variety of food, and the health services received are more limited [32].

The current study also concluded that subjects with a BMI \geq 25 have a 2.47 times greater risk of hypertension. This finding is consistent with previous studies, including a study in Poland, which states that having a BMI above average is a greater risk of hypertension [22]. Other studies, *i.e.*, Nagpur and South Asia, state the same conclusion [38]. A cohort study conducted in Thailand over four years showed that BMI was significantly associated with an increased risk of hypertension [26]. A study conducted among young adults in Tenwek Mission Hospital, Bomet County, Kenya, in 2018 stated that having a BMI \geq 25 had a 3.05 more significant risk of hypertension [39]. A Malaysian study stated that obese people have a 4.29 times risk of developing hypertension [40]. Likened to people with normal BMI, people with BMI \geq 25 are more likely to have metabolic disorders, such as fat metabolism disorder and insulin antibody disorder. Metabolic disorders can trigger cardiovascular disease, including hypertension [41]. Apart from other risk factors, BMI directly affects blood pressure. Nevertheless, the mechanism of BMI's influence on blood pressure is still unclear [42]. Several studies show that BMI is a critical and sensitive indicator for predicting high blood pressure [43, 44].

In developing countries, diabetes mellitus can be a significant factor in the development of cardiovascular disease [45]. This study showed that diabetes mellitus has a significant association with hypertension. People with diabetes mellitus are more at risk of developing hypertension [23, 26, 40, 46]. Hypertension and diabetes mellitus are interrelated because hypertension can also cause diabetes mellitus [47]. A previous study reported that the presence of diabetes mellitus puts a person at twice the risk of hypertension [48]. A prospective cohort study conducted over four years in a national Thai Cohort Study showed similar results, where diabetes mellitus was strongly associated with hypertension [26]. Diabetes mellitus is also a variable that affects hypertension, as demonstrated by a study conducted in Malaysia [40]. Research conducted in South Korea through The Fourth Korea National Health and Nutrition Examination Survey [KNHANES IV] states that hypertension is more prevalent in adults with diabetes mellitus in the general population and for both sexes [49]. Hypertension occurs due to complex metabolic disorders caused by diabetes mellitus. Increased circulating fluid volume and peripheral vascular resistance in patients with diabetes mellitus occur due to insulin resistance, which increases body fluid volume. This condition is closely related to increased blood pressure [50].

This study found that emotional mental disorder has a higher risk of getting hypertension. Psychologically, distress that occurs causes an increased risk of hypertension, 1.53 greater than a person without a mental disorder [51]. A study in China concluded that stress was significantly related to hypertension, where stressful conditions put a person 1.2 times more at risk of hypertension [52]. Emotional mental disorders, *e.g.*, stress, will trigger endothelial dysfunction that allows for an increase in blood pressure [53].

The present study also found that instant food consumption leads to hypertension condition. A prospective cohort study in Spanish revealed that high consumption of ultra-processed food creates a greater risk of hypertension [54]. An 8-year cohort study in Thailand showed that dietary patterns are associated with hypertension. An unhealthy diet includes consuming processed foods, including instant foods that are low in fiber, high in carbohydrates, saturated fat, and high in sodium [55]. Generally, the combinations contained in instant food are carbohydrates, fat, and salt [26]. Frequent consumption of instant food will trigger cardiometabolic, which causes an increase in blood pressure [56].

Since the study used secondary data regarding the data available in *RISKESDAS*, certain factors that influence the development of hypertension in young adults have yet to be identified. Future research is therefore anticipated to incorporate these characteristics, along with genetics and other factors, including modifiable factors and certain prevention efforts.

CONCLUSION

The prevalence of hypertension among young adults in this study was 4.1%, which showed lower numbers compared to other studies. This low prevalence might be due to a lack of awareness, low hypertension case detection, and underdiagnosis conditions, suggesting increasing early detection and screening efforts in the community.

Females, married, low-level education, unemployed, BMI \geq 25, diabetes mellitus, having an emotional mental disorder, and consumption of instant food altogether increase the risk of getting hypertension among young adults aged 26-35 years old.

AUTHOR'S CONTRIBUTION

L.R.P. and M.A.: Study conception and design; L.R.P. and A.I.F.: Data collection; L.R.P., M.A., A.A.N.: Analysis and interpretation of results. manuscript draft: LRP, MA, AIF, PK, and SAS. All authors reviewed the results and approved the final version of the manuscript.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This study was approved by the Ethical Review Board, Faculty of Medicine, Universitas Negeri Semarang, under the approval ID number 169/KEPK/FK/KLE/2024.

HUMAN AND ANIMAL RIGHTS

All human research procedures followed were in accordance with the ethical standards of the committee responsible for human experimentation (institutional and national), and with the Helsinki Declaration of 1975, as revised in 2013.

CONSENT FOR PUBLICATION

Informed consent was taken from the participants involved in this study.

AVAILABILITY OF DATA AND MATERIAL

The data supporting the findings of the article is available at <https://doi.org/10.6084/m9.figshare.27542946.v1>.

FUNDING

This work was financially supported by the Ministry of Education and Culture, Republic of Indonesia, and the Institute of Research and Community Service, Universitas Negeri Semarang [Grant number SP DIPA-023.17.1.690523/2024].

CONFLICT OF INTEREST

The author(s) declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

We also thank the National Institute for Health Development Policy, Ministry of Health, Republic Indonesia, for providing data from the Indonesian Basic Health Survey [Riset Kesehatan Dasar; RISKESDAS].

REFERENCES

- Unger T, Borghi C, Charchar F, et al. 2020 International society of hypertension global hypertension practice guidelines. *Hypertension* 2020; 75(6): 1334-57. <http://dx.doi.org/10.1161/HYPERTENSIONAHA.120.15026> PMID: 32370572
- World Health Organization. Hypertension. 2023. Available from: https://www.who.int/news-room/fact-sheets/detail/hypertension/?gad_source=1&gclid=Cj0KCQiAi_G5BhDXARISAN5SX7qWBFHjTIfkdpjBHCILhO_wjs0FnR78UKdDzLrel1eaOTznMFIIAbsaAlFueALw_wcB (accessed on 18-11-2024).
- Thabana MW, Grace JM. Selected non-communicable disease risk factors and mental health of south african durban-based refugees: A cross-sectional study. *Open Public Health J* 2024; 17(1): e18749445277768. <http://dx.doi.org/10.2174/0118749445277768240301072459>
- Azinar M, Azam M, Sugiharto S. Peer support group as a model for improving behavior of hypertension risk and complication control. *J Health Educ* 2019; 4(1): 17-21. <http://dx.doi.org/10.15294/jhe.v4i1.24192>
- Noncommunicable diseases. 2023. Available from: <https://www.who.int/data/gho/data/themes/noncommunicable-diseases#:~:text=They%20include%20tobacco%2C%20harmful%20use,blood%20sugar%20and%20raised%20cholesterol.> (accessed on 18-11-2024).
- The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. Bethesda (MD): National Heart, Lung, and Blood Institute (US) 2004.
- Amisi WG, Nelwan JE, Kolibu FK. The relationship between hypertension and the incidence of coronary heart disease in patients seeking treatment at the Central General Hospital, Prof. Dr R D Kandou Manado Kesmas 2018; 7(4): 1-7.
- Hypertension is the main cause of heart disease, kidney failure and stroke. 2021. Available from: https://p2ptm-kemkes-go-id.translate.google.com/informasi-p2ptm/hipertensi-penyebab-utama-penyakit-jantung-gagal-ginjal-dan-stroke?_x_tr_sl=id&_x_tr_tl=en&_x_tr_hl=en&_x_tr_pto=sc (accessed on 18-11-2024).
- Mills KT, Bundy JD, Kelly TN, et al. Global disparities of hypertension prevalence and control. *Circulation* 2016; 134(6): 441-50. <http://dx.doi.org/10.1161/CIRCULATIONAHA.115.018912> PMID: 27502908
- Johnson HM, Thorpe CT, Bartels CM, et al. Undiagnosed hypertension among young adults with regular primary care use. *J Hypertens* 2014; 32(1): 65-74. <http://dx.doi.org/10.1097/HJH.000000000000008> PMID: 24126711
- Iacobucci G. Adults at lowest risk are most likely to have undiagnosed hypertension, data show. *BMJ* 2023; 381: p976. <http://dx.doi.org/10.1136/bmj.p976> PMID: 37116940
- National Riset Kesehatan Report 2018 2018. Available from: https://repository-badankebijakan-kemkes-go-id.translate.google.com/eprint/3514/?_x_tr_sl=id&_x_tr_tl=en&_x_tr_hl=en&_x_tr_pto=sc (accessed on 18-11-2024).
- Depkes RI. Age Classification According to Category. Jakarta: Director General of Health Care 2009.
- Idaiani S, Suryaputri IY. The validity of the self-reporting questionnaire-20 for symptoms of depression: A sub-analysis of the national health survey in Indonesia. *Open Access Macedonian J Med Sci* 2022; 10: 1676-82.
- Loria CM, Liu K, Lewis CE, et al. Early adult risk factor levels and subsequent coronary artery calcification: The CARDIA Study. *J Am Coll Cardiol* 2007; 49(20): 2013-20. <http://dx.doi.org/10.1016/j.jacc.2007.03.009> PMID: 17512357
- De Venecia T, Lu M, Figueredo VM. Hypertension in young adults. *Postgrad Med* 2016; 128(2): 201-7. <http://dx.doi.org/10.1080/00325481.2016.1147927>
- Aggarwal R, Yeh RW, Joynt Maddox KE, Wadhwa RK. Cardiovascular risk factor prevalence, treatment, and control in US adults aged 20 to 44 years, 2009 to march 2020. *JAMA* 2023; 329(11): 899-909. <http://dx.doi.org/10.1001/jama.2023.2307> PMID: 36871237
- Ahammed B, Maniruzzaman M, Talukder A, Ferdausi F. Prevalence and risk factors of hypertension among young adults in albania. *High Blood Press Cardiovasc Prev* 2021; 28(1): 35-48. <http://dx.doi.org/10.1007/s40292-020-00419-5> PMID: 33113094
- Zahid Z, Tanvir H, Amjad A, Asim F, Sehar N, Asim F. Prevalence and associated factors of hypertension in young adults: A cross-sectional study. *Prof Med J* 2023; 31(1): 44-8. <http://dx.doi.org/10.29309/TPMJ/2024.31.01.7839>
- Krishnamoorthy Y, Rajaa S, Sulgante S, Chinnakali P, Jaswal N, Goel S. Prevalence of hypertension and determinants of treatment-seeking behaviour among adolescents and young adults in India: An analysis of NFHS-4. *J Public Health* 2023; 45(1): e48-56. <http://dx.doi.org/10.1093/pubmed/fdac006> PMID: 35233632
- Everett B, Zajacova A. Gender differences in hypertension and hypertension awareness among young adults. *Biodemogr Soc Biol* 2015; 61(1): 1-17. <http://dx.doi.org/10.1080/19485565.2014.929488> PMID: 25879259
- Wyszyńska J, Łuszczki E, Sobek G, Mazur A, Dereń K. Association and risk factors for hypertension and dyslipidemia in young adults from Poland. *Int J Environ Res Public Health* 2023; 20(2): 982. <http://dx.doi.org/10.3390/ijerph20020982> PMID: 36673736
- Singh S, Shankar R, Singh GP. Prevalence and associated risk factors of hypertension: a cross-sectional study in urban varanasi.

- Int J Hypertens 2017; 2017: 5491838.
<http://dx.doi.org/10.1155/2017/5491838>
- [24] Liu X, Xiang Z, Shi X, Schenck H, Yi X, Ni R. The risk factors of high blood pressure among young adults in the Tujia-Nationality Settlement of China. *Biomed Res Int* 2017; 8(1): 8315603.
<http://dx.doi.org/10.1155/2017/8315603>
- [25] Sabapathy K, Mwita FC, Dauya E, *et al.* Prevalence of hypertension and high-normal blood pressure among young adults in Zimbabwe: findings from a large, cross-sectional population-based survey. *Lancet Child Adolesc Health* 2024; 8(2): 101-11.
[http://dx.doi.org/10.1016/S2352-4642\(23\)00287-0](http://dx.doi.org/10.1016/S2352-4642(23)00287-0) PMID: 38070533
- [26] Thawornchaisit P, de Looze F, Reid CM, Seubsman S, Sleight AC, Chokhanapitak J. Health risk factors and the incidence of hypertension: 4-year prospective findings from a national cohort of 60 569 Thai Open University students. *BMJ Open* 2013; 3(6): e002826.
<http://dx.doi.org/10.1136/bmjopen-2013-002826> PMID: 23801711
- [27] Ramezankhani A, Azizi F, Hadaegh F. Associations of marital status with diabetes, hypertension, cardiovascular disease and all-cause mortality: A long term follow-up study. *PLoS One* 2019; 14(4): e0215593.
<http://dx.doi.org/10.1371/journal.pone.0215593> PMID: 31009512
- [28] Tiwari A, Datta BK, Haider MR, Jahan M. The role of child marriage and marital disruptions on hypertension in women - A nationally representative study from India. *SSM Popul Health* 2023; 22: 101409.
<http://dx.doi.org/10.1016/j.ssmph.2023.101409> PMID: 37132019
- [29] Sun K, Lin D, Li M, *et al.* Association of education levels with the risk of hypertension and hypertension control: A nationwide cohort study in Chinese adults. *J Epidemiol Community Health* 2022; 76(5): 451-7.
<http://dx.doi.org/10.1136/jech-2021-217006> PMID: 34996807
- [30] Pebrisiana P, Tambunan LN, Baringbing EP. Correlation of Characteristics with the Incident of Hypertension in Outpatients at RSUD Dr. Doris Sylvanus, Central Kalimantan Province. *Surya Medika J* 2022; 8(3): 176-86.
<http://dx.doi.org/10.33084/jsm.v8i3.4511>
- [31] Theofilou P. Associated factors with adherence level of elderly patients with hypertension abstract. *Open Public Health J* 2022; 15: 1-6.
<http://dx.doi.org/10.2174/18749445-v15-e221220-2022-136>
- [32] Rismadi K, Siagian A, Siregar FA. The influence of income and lifestyle on the incidence of hypertension in fishermen in the city of Medan. *Jurnal Health Sains* 2021; 2(3): 328-42.
<http://dx.doi.org/10.46799/jhs.v2i3.133>
- [33] Kato A, Kibone W, Okot J, Baruch Baluku J, Bongomin F. Self-reported hypertension and associated factors among adults in butambala district, central uganda: A community-based prevalence study. *Integr Blood Press Control* 2023; 16(November): 71-80.
<http://dx.doi.org/10.2147/IBPC.S434230> PMID: 37965566
- [34] Mohammed Nawi A, Mohammad Z, Jetly K, Abd Razak MA, Ramli NS, Wan Ibadullah WAH. The prevalence and risk factors of hypertension among the urban population in southeast asian countries: A systematic review and meta-analysis. *Int J Hypertens* 2021; 2021: 6657003.
<http://dx.doi.org/10.1155/2021/6657003>
- [35] Blok S, Haggensburg S, Collard D, *et al.* The association between socioeconomic status and prevalence, awareness, treatment and control of hypertension in different ethnic groups: The Healthy Life in an Urban Setting study. *J Hypertens* 2022; 40(5): 897-907.
<http://dx.doi.org/10.1097/HJH.0000000000003092> PMID: 35506279
- [36] Zheng C, Wang Z, Wang X, *et al.* Social determinants status and hypertension: A Nationwide Cross-sectional Study in China. *J Clin Hypertens (Greenwich)* 2020; 22(11): 2128-36.
<http://dx.doi.org/10.1111/jch.14023> PMID: 32882111
- [37] Patil CR, Sahoo DP, Dhole M, Kherde A, Inamdar A. Prevalence of hypertension and its associated risk factors in young adults attending a tertiary care institute of Nagpur: A cross sectional study. *Int J Community Med Public Health* 2017; 4(10): 3630.
<http://dx.doi.org/10.18203/2394-6040.ijcmph20174223>
- [38] Shahimi NH, Lim R, Mat S, Goh CH, Tan MP, Lim E. Association between mental illness and blood pressure variability: A systematic review. *Biomed Eng Online* 2022; 21(1): 19.
<http://dx.doi.org/10.1186/s12938-022-00985-w> PMID: 35313918
- [39] Ondimu DO, Kikui GM, Otieno WN. Risk factors for hypertension among young adults (18-35) years attending in Tenwek Mission Hospital, Bomet County, Kenya in 2018. *Pan Afr Med J* 2019; 33: 210.
<http://dx.doi.org/10.11604/pamj.2019.33.210.18407> PMID: 31692887
- [40] Zaki NAM, Ambak R, Othman F, *et al.* The prevalence of hypertension among Malaysian adults and its associated risk factors: Data from Malaysian Community Salt Study (MyCoSS). *J Health Popul Nutr* 2021; 40(S1): 8.
<http://dx.doi.org/10.1186/s41043-021-00237-y> PMID: 34059165
- [41] Tang N, Ma J, Tao R, *et al.* The effects of the interaction between BMI and dyslipidemia on hypertension in adults. *Sci Rep* 2022; 12(1): 927.
<http://dx.doi.org/10.1038/s41598-022-04968-8> PMID: 35042940
- [42] Landi F, Calvani R, Picca A, *et al.* Body mass index is strongly associated with hypertension: Results from the longevity check-up 7+ study. *Nutrients* 2018; 10(12): 1976.
<http://dx.doi.org/10.3390/nu10121976> PMID: 30551656
- [43] Wu X, Li B, Lin WQ, *et al.* The association between obesity indices and hypertension: Which index is the most notable indicator of hypertension in different age groups stratified by sex? *Clin Exp Hypertens* 2019; 41(4): 373-80.
<http://dx.doi.org/10.1080/10641963.2018.1489546> PMID: 30095294
- [44] Lee HA, Park H. Diet-related risk factors for incident hypertension during an 11-year follow-up: The Korean genome epidemiology study. *Nutrients* 2018; 10(8): 1077.
<http://dx.doi.org/10.3390/nu10081077> PMID: 30104485
- [45] calderón AM. Lifestyle risk factors associated with stroke in patients with type 2 Diabetes Mellitus: A case-control study abstract. *Open Public Health J* 2023; 16: 1-8.
- [46] Tsimihodimos V, Gonzalez-Villalpando C, Meigs JB, Ferrannini E. Hypertension and Diabetes Mellitus. *Hypertension* 2018; 71(3): 422-8.
<http://dx.doi.org/10.1161/HYPERTENSIONAHA.117.10546> PMID: 29335249
- [47] Azam M, Sakinah LF, Kartasurya MI, Fibriana AI, Minuljo TT, Aljunid SM. Prevalence and determinants of obesity among individuals with diabetes in Indonesia. *F1000 Res* 2023; 11: 1063.
- [48] Petrie JR, Guzik TJ, Touyz RM. Diabetes, hypertension, and cardiovascular disease: Clinical insights and vascular mechanisms. *Can J Cardiol* 2018; 34(5): 575-84.
<http://dx.doi.org/10.1016/j.cjca.2017.12.005> PMID: 29459239
- [49] Lee HS, Lee SS, Hwang IY, *et al.* Prevalence, awareness, treatment and control of hypertension in adults with diagnosed diabetes: The Fourth Korea National Health and Nutrition Examination Survey (KNHANES IV). *J Hum Hypertens* 2013; 27(6): 381-7.
<http://dx.doi.org/10.1038/jhh.2012.56> PMID: 23223084
- [50] Ohishi M. Hypertension with diabetes mellitus: physiology and pathology. *Hypertens Res* 2018; 41(6): 389-93.
<http://dx.doi.org/10.1038/s41440-018-0034-4> PMID: 29556093
- [51] Ojike N, Sowers JR, Seixas A, *et al.* Psychological distress and hypertension: Results from the national health interview survey for 2004-2013. *Cardiorenal Med* 2016; 6(3): 198-208.
<http://dx.doi.org/10.1159/000443933> PMID: 27275156
- [52] Hu B, Liu X, Yin S, Fan H, Feng F, Yuan J. Effects of psychological stress on hypertension in middle-aged Chinese: A cross-sectional study. *PLoS One* 2015; 10(6): e0129163.
<http://dx.doi.org/10.1371/journal.pone.0129163> PMID: 26043027
- [53] Munakata M. Clinical significance of stress-related increase in blood pressure: current evidence in office and out-of-office

settings. *Hypertens Res* 2018; 41(8): 553-69.

<http://dx.doi.org/10.1038/s41440-018-0053-1> PMID: 29808034

- [54] Mendonça RD, Lopes ACS, Pimenta AM, Gea A, Martinez-Gonzalez MA, Bes-Rastrollo M. Ultra-processed food consumption and the incidence of hypertension in a mediterranean cohort: The seguimiento universidad de navarra project. *Am J Hypertens* 2016; 30(4): hpw137.
<http://dx.doi.org/10.1093/ajh/hpw137> PMID: 27927627

[55] Shi Z, Papier K, Yiengprugsawan V, Kelly M, Seubsman S, Sleigh AC. Dietary patterns associated with hypertension risk among adults in Thailand: 8-year findings from the Thai Cohort Study. *Public Health Nutr* 2019; 22(2): 307-13.
<http://dx.doi.org/10.1017/S1368980018002203> PMID: 30187840

[56] Shin HJ, Cho E, Lee HJ, *et al.* Instant noodle intake and dietary patterns are associated with distinct cardiometabolic risk factors in Korea. *J Nutr* 2014; 144(8): 1247-55.
<http://dx.doi.org/10.3945/jn.113.188441> PMID: 24966409

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.