





Knowledge, Behaviors, and Treatment Associated with Sexually Transmitted Diseases: A One-Group Pretest-Posttest Study among Vietnamese Patients



Tam Thai Thanh Tran^{1,2} , Truc Thanh Pham³, Tin Trung Lam¹ , Tran Tran Ngoc Nguyen¹ , Nghia Nhu Nguyen¹ and Tin Hoang Nguyen^{1,2,*} 

¹Department of Physiology, Faculty of Medicine, Can Tho University of Medicine and Pharmacy, Can Tho City 900000, Vietnam

²Department of Functional Exploration, Can Tho University of Medicine and Pharmacy Hospital, Can Tho City 900000, Vietnam

³Le Van Thinh Hospital, Thu Duc City, Ho Chi Minh City 70000, Vietnam

Abstract:

Background: Sexually Transmitted Diseases (STDs), commonly referred to as venereal diseases, encompass several contagious ailments caused by a diverse array of microorganisms that vary in symptomatology, size, life cycle, and treatment resistance. The pathogens responsible for STDs include bacteria, viruses, fungi, and protozoa. Despite the diversity of STD pathogens, public understanding of transmission, prevention, and treatment remains limited. Additionally, changes in knowledge, behavior, and treatment practices following a non-pharmacological intervention remain controversial.

Objective: This research aimed to assess the STD status, knowledge, and behaviors of Vietnamese patients who visited the urology clinic at Le Van Thinh Hospital before and after both pharmacological and non-pharmacological interventions.

Materials and Methods: A one-group pretest-posttest design was used to assess the impact of two interventions on knowledge, behaviors, and treatment related to STDs. Participants (n=108) were evaluated 1 week and 3 months post-intervention. The interventions included pharmacological treatment, administered according to the Vietnam Ministry of Health guidelines, and a non-pharmacological intervention, including counseling, education, and awareness programs covering STD transmission, complications, prevention (e.g., condom use, health check-ups), and behavioral change. Data were collected through direct interviews, and SPSS was used for statistical analysis, with McNemar's non-parametric test employed to assess changes over time.

Results: Of all participants, 81.6% had gonorrhea. Remarkably, a complete lack of condom use was noted (100%). Before the intervention, 64.8% of participants were indicated to have a below-average or average level of understanding of STDs, and 100% participated in at least one risky behavior. Health education efforts led to a positive outcome, with 100% of participants showing good awareness. The majority of participants also showed reduced risky behaviors, with 52.8% using condoms. The therapeutic intervention achieved a 100% cure rate after 3 months, but limitations remained in the first week of treatment as three of the selected diseases demonstrated a treatment level of $\leq 25\%$.

Conclusions: The majority of the STD patients lacked knowledge and had a history of engaging in risky behaviors. The study demonstrated the effectiveness of current treatment protocols as well as the importance of education in increasing awareness and preventing risky behaviors.

Keywords: Venereal diseases, Condom, Educational intervention, Public health, Vietnam.

© 2025 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Received: December 23, 2024

Revised: March 04, 2025

Accepted: March 05, 2025

Published: June 02, 2025



Send Orders for Reprints to
reprints@benthamscience.net

* Address correspondence to this author at the Department of Physiology, Faculty of Medicine, Can Tho University of Medicine and Pharmacy, Department of Functional Exploration, Can Tho University of Medicine and Pharmacy Hospital, 179 Nguyen Van Cu Street, An Khanh Ward, Ninh Kieu District, Can Tho City 900000, Vietnam; E-mail: nhtin@ctump.edu.vn

Cite as: Tran T, Pham T, Lam T, Tran Ngoc Nguyen T, Nguyen N, Nguyen T. Knowledge, Behaviors, and Treatment Associated with Sexually Transmitted Diseases: A One-Group Pretest-Posttest Study among Vietnamese Patients. Open Public Health J, 2025; 18: e18749445348558. <http://dx.doi.org/10.2174/0118749445348558250428073432>

1. INTRODUCTION

Sexually Transmitted Diseases (STDs) are widespread infections, primarily affecting young people. The World Health Organization (WHO) reports millions of new cases daily, but the true burden remains uncertain due to asymptomatic cases, limited diagnostics, and inadequate surveillance systems worldwide [1, 2]. STDs impact both physical and mental health, with a risk of vertical transmission during pregnancy. Syphilis can cause stillbirth and neonatal complications, while human papillomavirus may lead to cervical cancer. Gonorrhea and chlamydia increase the risk of pelvic inflammatory disease, infertility, and premature birth [1, 3]. Owing to their prevalence and associated problems, STDs substantially impact the quality of life and mortality rates. Thus, the issue has transcended the strictly medical context, entering the domain of urgent societal responsibilities [2, 4]. In recent years, the prevalence of STDs has risen markedly due to the premature initiation of sexual activity, frequent partner changes, disregard for condom usage, insufficient sex education, and self-medication, among other factors [5, 6].

The decreasing age at which sexual intercourse begins has been identified as a contributing factor to the rise in STD incidence [5, 7]. Data from several European nations indicate that the average age of initial sexual intercourse has declined over the past 30 years, with a growing percentage of adolescents reporting sexual activity before the age of 16 years [8]. The premature initiation of sexual activity elevates the likelihood of engaging with several sexual partners and heightens the risk of acquiring a sexually transmitted infection [9, 10]. The risk is elevated for female teenagers due to their underdeveloped cervical anatomy, rendering them particularly susceptible to specific STDs [10]. Adolescents often avoid condoms due to inexperience or discomfort and focus solely on pregnancy prevention, reducing STD protection [9]. Moreover, numerous adolescents do not consider themselves at risk of acquiring STDs [11]. As the number of STD clinics has diminished, patients have increasingly sought care for STDs at general care clinics, emergency rooms, and family planning clinics [12]. General care physicians play a key role in STD detection, as many cases are asymptomatic. Primary care clinics diagnose up to 50% of reported STDs [7, 10]. In 2022, 71% to 80% of STD cases were recorded from non-STD clinics [13]. Expanded health insurance led to a 100% increase in primary care

visits for STD care, while visits to STD clinics dropped by 20%. The rise was mainly driven by more women seeking STD treatment [14, 15]. Notwithstanding these changes in the venues for STD service delivery, publicly financed STD clinics remain a crucial safety net for uninsured individuals and other vulnerable groups seeking care [11].

Managing STDs is complicated due to the prevalence of asymptomatic or moderate infections, which complicate detection. The variety of infectious comorbidities and social stigma also hinder patient-centered care; thus, multidisciplinary management is crucial [10, 12]. The WHO promotes STD education to reduce the disease burden. A study found that young students were mostly aware of Human Immunodeficiency Virus/Acquired Immunodeficiency Syndrome (HIV/AIDS), with limited knowledge of other STDs, mainly acquired from the Internet and media [9, 11, 16]. Many people, especially in developing countries, underestimate STD risk and lack awareness. Therefore, understanding symptoms and prevention is crucial to reducing transmission and infection rates [5, 7].

The incidence of STDs in metropolitan regions and megacities is rising. Previous studies have elucidated a correlation between the proliferation of STDs and extensive urbanization and migration. Moreover, in underdeveloped nations, STDs have been demonstrated to adversely affect public health, significantly impacting morbidity and mortality rates. Research on STDs in Vietnam is limited; therefore, this study aimed to assess the STD status, knowledge, and behaviors of participants and to evaluate changes in their knowledge and behaviors before and after intervention.

2. MATERIALS AND METHODS

2.1. Study Design and Setting

Our study involved a one-group, pretest-posttest design and was conducted on 108 patients who visited the urology clinic at Le Van Thinh Hospital, Thu Duc City, for STD testing and treatment. Participants were invited to complete 20-minute face-to-face interviews using structured questionnaires administered by well-trained researchers.

2.2. Sample size

The Following Formula was Used:

$$n = Z^2_{1-\alpha/2} \frac{(1-\rho)}{\rho(\varepsilon)^2} \quad [17]$$

In this formula, n is the sample size; $Z_{1-\alpha/2} = 1.96$ is the 95% confidence interval; $\rho = 90.8\%$ is the proportion of patients who understand STDs [5]; and $\varepsilon = 0.062$ is a relative precision, calculated as 6.2% of ρ . Substituting the relevant values into the above formula, the minimum sample size for the study was 102 patients; in total, 108 patients were included.

2.3. Pilot study

A pilot study was conducted to assess the suitability of the questionnaire, evaluate the data collection process, and estimate initial parameters in preparation for the main study. The pilot study was conducted on a small sample of 20 individuals representing the target population, using direct interviews as the data collection method. This process included administering the survey, assessing participants' feedback, and evaluating input on the clarity of the questions, completion time, and any issues encountered during implementation. The results helped identify errors or limitations in the questionnaire design and provided preliminary data for adjusting the sample size for the main study.

2.4. Interventions

Pharmacological interventions were administered based on the identified causative agents, strictly adhering to the treatment regimens outlined in the general guidelines of the Ministry of Health (Vietnam) to ensure standardized and effective care. Additionally, non-pharmacological interventions focused on educating the study subjects about their conditions and raising awareness of STDs were administered. These educational efforts included providing comprehensive information on the nature, transmission, and potential complications of STDs, as well as equipping individuals with knowledge of preventive measures, such as safe sexual practices, condom use, and regular health check-ups. To reinforce awareness, counseling sessions were conducted, and educational materials were disseminated to foster long-term behavioral changes that could contribute to reducing disease transmission within the community.

2.5. Data collection

The questionnaire was developed in Vietnamese with additional revisions suitable for Vietnamese patients (Supplementary Material). It was divided into six parts (eight questions about socioeconomic characteristics, 10 questions about the participants' STD status and risky behaviors, 18 questions about their awareness of STDs before and after intervention, four questions about treatment results after a week, and four questions about treatment results after 3 months, and 10 questions about behaviors before and after intervention) and comprised multiple-choice questions (Fig. 1).

2.6. Questionnaire Validity

To validate the questionnaire on STD-related knowledge,

behaviors, and treatment in a one-group pre-post study, key statistical methods were used to ensure reliability. Content validity was assessed by experts using the content validity index, requiring a score of ≥ 0.80 . Criterion validity was determined by comparing questionnaire scores to a gold standard using Pearson's correlation, with an acceptable $r > 0.7$.

2.7. Reliability Analysis

Reliability analysis ensured that the questionnaire on STD knowledge, behaviors, and treatment consistently measured what it intended. Internal consistency was assessed using Cronbach's alpha (≥ 0.7), while test-retest reliability evaluated stability over time with the intraclass correlation coefficient (≥ 0.75). Together, these methods confirmed the questionnaire's dependability for research and public health.

2.8. Data Analysis

Data from the collected forms were entered and processed using SPSS 20.0 software and medical statistics methods. Descriptive statistics (frequencies and percentages) were used for the following qualitative variables: gender, age group, marital status, occupation, education, income, place of residence, history of STDs, STD status, knowledge about STDs, risk behaviors for STDs, and treatment outcomes. Comparative analysis of changes in knowledge and behaviors about STDs before and after intervention was performed using McNemar's non-parametric test.

2.9. Ethical Considerations

All study participants provided informed consent before data collection, having fully understood the purpose, risks, and benefits of their participation. To protect participants' personal information, strict confidentiality measures were implemented. All collected data (Supplementary Material) were anonymized, securely stored in encrypted databases, and accessible only to authorized researchers. Identifiable information was removed or coded to prevent any link to individual participants. Additionally, no personal data were shared or disclosed beyond the research team. Participants had the right to withdraw at any stage without consequences, ensuring that their autonomy and privacy were fully respected.

3. RESULTS

3.1. Sociodemographic Characteristics of the Participants

Table 1 shows that most participants were men aged 20–49, unmarried, and residing in urban areas. They worked in various professions, primarily as workers and office staff, most of whom had an education level of high school or higher and a high income, in contrast to those with lower income and educational attainment. Furthermore, most participants reported no history of STDs, while a small proportion had previously contracted gonorrhoea.




Fig. (1). Study flowchart.

Table 1. Sociodemographic characteristics of the participants.

Characteristics		Frequency	Percentage (%)
Age	< 20	5	4.6
	20-29	42	38.9
	30-39	34	31.5
	40-49	18	16.7
	50-59	6	5.6
	> 60	3	2.7
Gender	Male	105	97.2
	Female	3	2.8
Marital status	Unmarried	53	49.1
	Married	31	28.7
	Divorced/widowed	24	22.2
Education level	Elementary	9	8.3
	Secondary	14	13.0
	High school	43	39.8
	College/university/postgraduate	42	38.9

Characteristics		Frequency	Percentage (%)
Occupation	Office staff	28	25.9
	Worker	27	25.0
	Driver	15	13.9
	Trader	16	14.8
	Student	12	11.1
	Farmer	5	4.6
	Others	5	4.6
Income	High-income	78	72.2
	Upper-income	16	14.8
	Middle-income	7	6.5
	Low-income	7	6.5
Residence	Urban	76	70.4
	Rural	32	29.6
History of STDs	No	94	87.0
	Yes (gonorrhoea)	14	13.0

3.2. STD Status and Risk Behaviors of the Participants

As summarized in Table 2, most participants had already been diagnosed with STDs (primarily gonorrhea) at enrollment. The most common risk behavior was a lack of condom use, followed by having sex with sex workers. In contrast, only a few were positive for HIV or syphilis. Additionally, men who have sex with men (MSM) represented a small proportion of the study group.

3.3. Treatment Results

The treatment efficacy observed after the first week

was relatively low, with three of the selected diseases demonstrating a treatment level of $\leq 25\%$, no recovery among syphilis cases, and only chlamydia reaching 100% efficacy. However, after 3 months, all patients achieved full recovery. These results are presented in Table 3.

3.4. Post-intervention Results

Table 4 shows positive changes regarding the participants' involvement in reducing risk factors and enhancing preventive measures, including increased condom use and reduced sex with sex workers. However, only a minority completely avoided all risky behaviors post-intervention.

Table 2. Sexually Transmitted Disease (STD) status and risk behaviors of the participants.

Characteristics		Frequency	Percentage (%)
STD status	Gonorrhoea	93	86.1
	Chlamydia	7	6.4
	Genital warts	4	3.7
	Syphilis	2	1.9
	HIV/AIDS	2	1.9
Risk behaviors	Not using condoms	108	100
	Having sex with sex workers	89	82.4
	Multiple sexual partners (> 2)	74	68.5
	Stimulant usage during sexual activities	54	50.0
	Men who have sex with men	11	10.2

Table 3. Treatment results after a week and 3 months.

STDs	Cases	Treatment Success after a Week		Treatment Success After 3 Months		*p-value
		Frequency	Percentage (%)	Frequency	Percentage (%)	
Gonorrhoea	93	5	5.4	93	100	< 0.001
Syphilis	2	0	0	2	100	0.157
Chlamydia	7	7	100	7	100	1
Genital warts	4	1	25	4	100	0.0832

*The McNemar test was employed to compare the percentages pre- and post-intervention.

Table 4. Participants' behaviors regarding STDs before and after the intervention.

Characteristics	Pre-intervention		Post-intervention		*p-value
	Frequency	Percentage (%)	Frequency	Percentage (%)	
Sexual partners					
1 person	34	31.5	42	38.9	< 0.001
At least 2 people	74	68.5	66	61.1	
Having sex with sex workers					
Yes	89	82.4	45	41.7	< 0.001
No	19	17.6	63	58.3	
Using condoms					
Yes	0	0.0	57	52.8	< 0.001
No	108	100	51	47.2	
Men who have sex with men					
Yes	11	10.2	8	7.4	< 0.001
No	97	89.8	100	92.6	
General comparison of risk behavior changes					
Yes	108	100	94	87.0	< 0.001
No	0	0	14	13.0	

*The McNemar test was employed to compare the percentages pre- and post-intervention.

3.5. Participants' Knowledge of STDs Before and After Intervention

As presented in Table 5, most participants had a medium level of knowledge about STDs, especially STD symptoms, common risk behaviors, and transmission routes, such as

having sexual intercourse without con-doms. However, few were aware of parasitic causative agents, transmission through unsafe blood transfusions or needle sharing, and complications. After the intervention, the participants achieved excellent knowledge levels compared to before, particularly on STD complications and transmission routes.

Table 5. Participants' knowledge about STDs before and after the intervention.

Characteristics	Pre-intervention		Post-intervention		*p-value
	Frequency	Percentage (%)	Frequency	Percentage (%)	
Knowledge about common STDs					
Gonorrhoea	101	93.5	107	99.1	0.003
Syphilis	88	81.5	107	99.1	< 0.001
Genital warts	83	76.6	105	97.2	< 0.001
HIV/AIDS	76	70.4	108	100	< 0.001
Genital herpes	23	21.3	83	76.9	< 0.001
Chlamydia	13	12.0	55	50.9	< 0.001
<i>Trichuris trichiura</i>	7	6.5	31	28.7	< 0.001
Knowledge of STD pathogens					
Virus	84	77.8	105	97.2	< 0.001
Bacteria	75	69.4	105	97.2	< 0.001
<i>Trichomonas, Candida, Sarcoptes scabiei, Pthirus pubis</i>	20	18.5	85	78.7	< 0.001
Knowledge of STD symptoms					
List at least three correct symptoms	93	86.1	108	100	< 0.001
List fewer than three symptoms	15	13.9	0	0	< 0.001
Knowledge of STD risk behaviors					
Not using condoms	99	91.7	108	100	< 0.001
Having sex with sex workers	89	82.4	106	98.2	< 0.001
Multiple sexual partners	59	54.6	104	96.3	< 0.001
Unsafe blood transfusion	33	30.6	79	73.2	< 0.001
Knowledge of STD transmission					
Sexual activities	102	94.4	105	97.2	< 0.001
Blood transfusion	48	44.4	101	93.5	< 0.01
From mother to child	46	42.6	92	85.2	< 0.001
Sharing a needle pump	38	35.2	61	87.0	< 0.001

Characteristics	Pre-intervention		Post-intervention		<i>*p-value</i>
Knowledge of STD complications					
List at least three complications	48	44.4	107	99.1	< 0.001
List fewer than three complications	60	55.6	1	0.9	< 0.001
General comparison of knowledge changes					
Bad	31	28.7	0	0	< 0.001
Medium	39	36.1	0	0	< 0.001
Good	31	28.7	12	11.1	< 0.001
Excellent	7	6.5	96	88.9	< 0.001

*The McNemar test was employed to compare the percentages pre- and post-intervention.

4. DISCUSSION

4.1. Sociodemographic Characteristics

The characteristics of our study cohort were consistent with those typically observed among patients visiting the urology department of Le Van Thinh Hospital. Existing research also indicates that men are significantly more likely than women to seek clinical care for STDs. Similarly, it has been found that women tend to delay seeking help for sexual health issues longer than men—on average, 41 days compared to 16 days [18]. These gender differences may be attributed to the fact that sexual health remains a sensitive topic for many women. In numerous Asian cultures, persistent cultural stigma continues to discourage open discussion of STDs among women.

The large number of participants within the age range of 20–29 stems from the fact that this final stage of sexual development, coupled with curiosity, leads individuals to have a strong tendency to engage in sexual activities. Moreover, individuals want to prove their maturity and experience in the field of sexuality. Therefore, many of them seek out sexual activities. The above speculation is reinforced by a study conducted by Hanoi Medical University on 873 male patients who had STDs. The incidence of STDs in the age group of 20–30 years accounted for 44.33% of cases, and this age group was determined to be the one with the highest levels of physiological activity and sexual impulses that tended to seek out sexual partners to satisfy physiological needs [19]. In addition, the WHO estimated that, in addition to AIDS, if only four treatable diseases are taken into account (trichomonas, chlamydia, polio, and syphilis), more than 340 million new cases occur globally among 15–49-year-olds, underscoring this high-risk demographic [20]. Due to the complex nature of sexual life and the high rates of STDs reported previously, further research is required on this at-risk age group.

The fact that the majority of study participants were single and not bound by marriage and romantic relationships led them to engage in more vigorous sexual activity compared to those who were married or in a romantic relationship. Similarly, in a study conducted in 2022 on male subjects with STDs at the Hanoi Medical University Hospital, unmarried participants accounted for almost half of the participants, at 46.09% [21]. Conversely, a study on patients who visited the Suraksha Clinic under the Dermatology Department in Lady Hardinge Medical

College, New Delhi state, India, from 2003 to 2019 found that the prevalence of STDs was higher in married persons (81.7%) than in unmarried people (17.9%) [22], highlighting how cultural norms shape epidemiological patterns.

We observed a relationship between academic qualifications and the incidence of STDs because the majority of individuals with STDs had an education level of high school or higher. The high STD infection rate among individuals with a high school education or above contradicts initial assumptions, as these individuals are typically well-equipped with knowledge about STDs. However, these results show some similarities with a study revealing that academic qualification from the high school level onward accounted for 88.2% of 622 patients examined at the Vietnam National Hospital of Dermatology and Venereology from September to November 2018 [23]. In a 2021 study on drug use and health conducted by the United States Intramural Research Program at the National Institutes of Health, approximately 32.43% of participants held a college degree or higher [24]. Interestingly, the rising rates of STD infections among individuals with higher education levels may reflect a lack of emphasis on sexual health education and a general indifference toward preventive measures. This may be particularly true in educational systems where sex education is not prioritized alongside core subjects, such as English and mathematics. Further research is needed to better understand the underlying factors contributing to this concerning trend among well-educated populations.

Occupation and accommodation status also influenced the study results. The characteristics of an individual's living environment and educational background can shape their cognitive processes and health-related behaviors. In our study, most participants were laborers or students—groups that often have limited exposure to public health education programs and possess less knowledge about sexual health and safety. Moreover, rapid urbanization and the pursuit of economic opportunities have led to population growth in major cities, particularly in Ho Chi Minh City. This demographic shift complicates epidemiological control and contributes to rising rates of STDs in urban areas, as also reported in Smith's study on the Asia-Pacific region [22]. Additionally, higher income levels may enable engagement in riskier behaviors, such as substance use and relationships with

sex workers, further increasing STD risk. However, it is equally important to consider rural and low-income populations, who often lack access to healthcare services and may, therefore, be underrepresented in official statistics. These findings highlight the need for further research into the relationships between place of residence, income level, and STD prevalence.

4.2. Incidence of STDs and Risk Behaviors of Patients with STDs

4.2.1. Incidence of STDs

Since this study was conducted during the fourth phase of the coronavirus disease 2019 (COVID-19) pandemic, when Vietnam implemented social distancing measures, along with the general apprehension of the public and patients regarding the risk of COVID-19 transmission, the capacity for screening tests and the identification of individuals who have been or are at risk of contracting STDs was limited. On the other hand, prioritizing resources for preventing the COVID-19 pandemic may have led to reduced testing and diagnosis of at-risk individuals. Therefore, according to the researchers' estimates, the number of people with STDs may be much higher.

Furthermore, a study in Vietnam reported a rapid increase in the incidence of sexually transmitted infections over the past decade, reaching 17.3% in 2017, with an estimated 200,000 new cases annually [25]. Similarly, a major survey conducted in the United States in 2018 identified up to 32 million men living with an STD [26]. This substantial pre-pandemic burden, combined with the current lack of widespread screening, raises concerns about a potential significant rise in sexual health issues following the COVID-19 pandemic.

The majority of patients in our study were diagnosed with gonorrhea, making it the most commonly identified sexually transmitted disease in our sample. This distribution aligns with the prevalence patterns reported in previous studies. For example, in Spain, the incidence of gonorrhea has been increasing since the early 2000s [27]. Additionally, a WHO study reported over 82 million cases of gonorrhea globally in 2021 alone [28]. The higher prevalence of gonorrhea compared to other STDs may be attributed to its high transmissibility and the growing antimicrobial resistance of *Neisseria gonorrhoeae* to all therapeutic antibiotics [29], which reduces treatment efficacy [30]. Furthermore, many individuals with gonorrhea, especially women, often exhibit no obvious symptoms or only mild signs, such as unusual vaginal discharge, pain during urination, and lower abdominal pain, symptoms that are commonly mistaken for other urinary tract infections.

Although HIV/AIDS was recorded at a low rate in our study, it remains a significant public health concern due to challenges in controlling its spread, particularly driven by unsafe sexual practices and increased injection drug use. Notably, sexual transmission of HIV among men who have sex with men (MSM) has risen rapidly in recent years.

According to the Department of Prevention, Anti-HIV/AIDS (Vietnam Ministry of Health), the primary modes of HIV transmission in Vietnam are needle sharing among people who inject drugs and sexual contact with sex workers. However, sexual transmission, especially within the MSM population, has been on the rise [31]. The average HIV infection rate among MSM increased from 6.7% in 2014 to 12.2% in 2017, reaching 13.3% in 2020. In some regions, more than 50% of newly diagnosed HIV cases were among MSM. Despite this, in Ben Tre province in southern Vietnam, only 79% of MSM had undergone HIV testing in 2018 [32]. Sexual activity among MSM is recognized as a major risk factor driving the HIV epidemic both in Vietnam [7] and globally [20]. Therefore, further research focused on the sexual health needs of MSM populations is urgently needed.

4.2.2. Risk Behaviors

This study included several questions targeting behaviors known to be risk factors for disease transmission, particularly unsafe sexual practices extensively documented in public health reports, media, and legal texts [32]. Alarming, all participants reported not using condoms, a finding that raises significant concerns. The absence of condom use eliminates the physical barrier that prevents direct contact between genitalia and bodily fluids, thereby facilitating pathogen transmission. Notably, many studies have consistently reported low rates of condom use among young people [32].

In addition to inconsistent condom use, the relatively high proportion of individuals engaging in sexual activity with multiple partners serves as a further warning sign. Having multiple sexual partners increases the likelihood of pathogen exposure and complicates efforts to monitor and manage individuals' health status. For men, each additional sexual partner increases the risk of contracting STDs by approximately 6% [33]. Moreover, sexual relations with multiple partners contribute to cross-transmission within the community, particularly among sex workers, who frequently have unprotected sex with multiple clients. This makes them highly vulnerable to STDs and accelerates disease spread. Reported STD rates among female sex workers range from approximately 33% to 50% [34].

Alcohol and drug use before and during sexual activity remain important contributing risk factors. In Brookmeyer's study, the proportion of stimulant users was notably high at 86.4%, with a statistically significant p-value of less than 0.001 [35]. This finding strongly suggests a link between stimulant use and an increased risk of STDs. One possible explanation for this association is that stimulant use impairs self-control and decision-making abilities. Individuals under the influence of stimulants often experience heightened impulsivity, reduced inhibitions, and impaired judgment. As a result, they may engage in risky sexual behaviors, such as unprotected sex or multiple sexual encounters, without fully considering the potential consequences. The

stimulating effects of these substances can lead to reckless and spontaneous decisions, further increasing the likelihood of STD transmission [36].

Besides these widely known factors, homosexual relationships are also notable, even though they have not yet emerged as a predominant source of STDs in the general community. This population is considered at elevated risk due to the biological vulnerability associated with unsafe sexual practices, such as anal sex and oral sex, leading to higher transmission rates of STDs [7]. Additionally, stigma and discrimination toward homosexuality prevent such individuals from accessing healthcare services, resulting in many MSM not receiving timely testing and treatment. A 2014 report from the United States Centers for Disease Control and Prevention showed that of the total of 91% of syphilis cases in men, 83% were among MSM. Moreover, MSM accounted for up to 51% of reported HIV cases across 26 states in the United States [37]. Further research is essential to develop solutions aimed at reducing the STD rate among this group.

4.3. Participants' Knowledge of STDs

Despite ongoing efforts to raise public awareness about STDs, our findings identified a relatively low level of understanding. Such limited knowledge may contribute to increased transmission of STDs. Other studies have also noted that a lack of understanding regarding STDs can hinder disease prevention efforts [38, 39]. Although our results indicated that most participants demonstrated knowledge of common STDs, a significant proportion lacked understanding of less common STDs, such as chlamydia, which are becoming potential threats due to a recent rise in incidence [35].

Viruses and bacteria were widely recognized as causative agents. In contrast, parasites were less commonly identified, likely because they are rarely mentioned in the media and may be easily confused with viruses and bacteria due to a lack of knowledge. Interviews with rural Vietnamese women on their knowledge of STDs revealed that 17% of participants classified *Phthirus pubis* as an STD pathogen, while 52% and 34%, respectively, classified it as bacterial candida and vaginitis [40]. Such misconceptions pose significant challenges to disease prevention and treatment, particularly in underserved areas with limited access to diagnostic equipment.

STD symptoms were widely recognized in this study. In contrast, Nguyen Huu Sau reported that patients could only list some symptoms of STDs, including urethral ulcers (53.5%), genital redness (49.6%), and genital ulcers (48.8%) [41]. Moreover, in Mansor's research, about 90% of participants stated that STD patients may not show any symptoms or signs [42]. These previous studies differ from our research in the level of understanding regarding STD symptoms. This difference may stem from variations in sample size and the level of knowledge about STDs between residential areas.

Regarding STD complications, a relatively low percentage of participants were aware of the potential

consequences. This finding aligns with a study conducted among college students in India, where fewer than 50% of respondents had knowledge about STD complications [43]. The lack of awareness, particularly among individuals with higher education levels, may be attributed to limitations in sexual health education and psychological reluctance to openly address sexual health issues. Furthermore, public health efforts have often focused on disseminating information about a limited number of common STDs, resulting in insufficient knowledge about other infections. For example, previous studies in Vietnam have demonstrated low levels of accurate knowledge concerning complications associated with chlamydia, such as male infertility, pelvic inflammatory disease, and ectopic pregnancy [44], largely because chlamydia remains relatively unknown in the general population.

Not using condoms remains a significant and well-recognized risk factor for STD transmission, as widely highlighted in both research and media outlets. A study in Brazil on military recruits also reached a similar conclusion, with 97.2% of participants believing that not using condoms during sexual intercourse increases the risk of contracting STDs [45]. Although participants were aware of the potential dangers of risky behaviors in Mansor's study, they could not name the two most effective preventive measures for STDs, that is, having sexual relations with one partner and using condoms during sexual intercourse [46]. The aforementioned study reflects the reality that important methods for preventing STDs are not well known, even though there is considerable knowledge about risky behaviors.

Apart from sexual intercourse being widely recognized as a common transmission route, only a few participants were unaware of the remaining transmission pathways, such as blood transfusion, mother-to-child, and needle sharing, which originally received little attention. This raises concerns about the lack of preventive measures for unpopular transmission pathways, particularly regarding needle sharing, due to its relevance to drug use, which is among the main drivers of STD transmission, especially HIV/AIDS [35].

4.4. Treatment Success, Risk Factors, and Participants' Knowledge of STDs After the Intervention

4.4.1. Treatment Success

The clinical success rate for gonorrhea after one week of treatment in our study was comparable to rates reported in other research. For instance, Vanessa and colleagues reviewed medical records of 291 patients infected with *Neisseria gonorrhoeae* who were treated with cefixime between May 1, 2010, and April 30, 2011, at a sexual health hospital in Toronto, Canada, reporting a success rate of 6.77% [47]. Similar success rates for cefixime and azithromycin in treating gonorrhea have been observed in studies from the United States and Europe [48, 49]. Although all patients in our study achieved full recovery within three months, the initial

failure rate after one week raises concerns about increasing antimicrobial resistance, a problem highlighted by other researchers [50]. Continued investigation is needed to develop more effective treatment strategies.

Among five patients treated for genital warts, four underwent minor surgical removal procedures, while one received trichloroacetic acid therapy [51]. However, after one week, two patients who had surgery and the patient treated with trichloroacetic acid continued to exhibit symptoms. This finding is noteworthy, as many reports indicate treatment effectiveness rates of up to 86% [52]. It is important to consider, however, that the small sample size in our study may limit the reliability of these results and could contribute to discrepancies in conclusions regarding treatment efficacy.

Regarding chlamydia, complete success was achieved when the Department of Health treatment scheme was applied. Similar results were noted in MSM subjects with rectal chlamydia, where 100% of the cases in the doxycycline treatment group recovered, while only 74% of the subjects in the azithromycin treatment group showed positive results [51]. However, azithromycin is highly regarded for treating patients with erratic healthcare-seeking behaviors, poor treatment compliance, or unpredictable follow-up due to its single-dose regimen [52]. Therefore, the appropriate treatment plan should be decided based on the patient's lifestyle and work characteristics.

Syphilis treatment is based on benzathine penicillin 2.4 million units, with the appropriate injection dosage depending on the disease stage [39]. Alternative therapies are considered in cases of low treatment effectiveness. According to the European guideline on the management of syphilis, intramuscular procaine penicillin for 10-14 days is the second-line option, along with other alternatives, such as doxycycline or azithromycin in cases of coagulopathy, penicillin allergy, or refusal of injectable treatment [34]. However, in clinical practice, the therapeutic efficacy of these alternative therapies is often limited, with reports of drug resistance to azithromycin and a higher failure rate for doxycycline compared to benzathine penicillin [30, 31, 35]. Therefore, benzathine penicillin remains the most trusted choice. Given the decreasing effectiveness of treatment, further research is needed to identify appropriate therapies for syphilis.

4.4.2. Risk Behaviors and Participants' Knowledge After the Intervention

The results of this quasi-experimental study indicated that the STD risk prevention intervention administered to patients at Le Van Thinh Hospital increased and sustained knowledge about STDs as well as promoted safer sexual behaviors, with monitoring indicating a reduction in STD prevalence. Our research provides evidence that effective educational interventions lead to gains in knowledge about STDs and shows that behavioral intentions can be modified toward "safer" sexual activities.

Through knowledge-sharing consultations, brochure distribution, and instructions on condom use, successive surveys demonstrated a significant increase in STD-

related knowledge throughout the study. The proportion of participants with "excellent" knowledge rose from 6.5% at baseline to 88.9%, while the remaining 11.1% achieved a "good" level of understanding. A statistically significant improvement was also observed in participants' intention to always use condoms. This improvement in condom use is a crucial foundation for preventing and reducing the incidence of STDs. Additionally, the positive changes in STD knowledge contribute to strengthening prevention and control efforts, particularly among adolescents, and may help reduce premarital sexual activity [52]. However, our findings also revealed minimal changes in awareness of other risk factors, such as unsafe blood transfusions and needle sharing. Further research targeting these specific areas is necessary to enhance public understanding.

Overall, this study demonstrates the benefits of STD prevention through comprehensive sex education and educational reform, including the integration of information on STDs, risk behaviors, transmission routes, and less common pathological features. These efforts contribute to improved knowledge, the promotion of safer sexual practices, and a reduction in risky behaviors, particularly among highly educated individuals. The findings may also support the broader use of technology in STD-related education. Experts from the European Society for Sexual Medicine have emphasized that digital sex education, particularly through smartphone applications, is emerging as a new trend in sexual health education [49], offering rapid and accurate access to information and enabling better epidemiological monitoring of target populations.

5. LIMITATIONS

This study is subject to several limitations that should be considered when interpreting the findings. Since the research was conducted during the fourth phase of the COVID-19 pandemic, when Vietnam had implemented a lockdown policy, fewer patients could participate in the study, which limited the sample size. This limits generalizability, reduces accuracy, and decreases the statistical power of the study, as well as increases the margin of error. Additionally, the overreliance on binary (yes/no) questions may have introduced response bias, thereby affecting the objectivity and validity of the data collected. Therefore, further research is necessary to address these weaknesses.

CONCLUSION

Through this research, STD-related epidemiology, knowledge, and behaviors were recorded and summarized. By assessing STD status alongside social and lifestyle characteristics, the study provided insight into the behavioral patterns and knowledge levels of the cohort. Approximately 70% of participants demonstrated below-average knowledge. While most participants were aware of symptoms, risk behaviors, and common transmission routes (such as sexual intercourse and lack of condom use), their understanding of complications, additional risk factors, and less common transmission routes (such as parasitic infections and unsafe blood transfusions) was

limited. Furthermore, participation in risky behaviors, particularly the universal lack of condom use, contributed to the 100% STD infection rate observed among participants prior to the intervention.

After the intervention, participants demonstrated positive changes, including improved knowledge of causative agents, complications, and transmission pathways, as well as a reduction in risk behaviors. More than half of the participants reported condom use, and there was a noticeable decline in high-risk sexual behaviors among MSM individuals.

We also evaluated the effectiveness of current treatment protocols, which showed positive results after 3 months of treatment but had limited efficacy in the first week. Taken together, this one-group pretest-posttest study provides an objective perspective and provides a basis for designing measures to prevent, mitigate, and treat STDs.

AUTHORS' CONTRIBUTIONS

The authors confirm their contributions to the paper: study conception and design: Tam T.T.T. and Truc T.P. were responsible for the study conception and design, as well as data collection. Tin H.N., Tin T.L., and Tran T.N.N. conducted the analysis and interpretation of the results. The initial draft of the manuscript was written by Nghia N.N., Tin T.L., Tran T.N.N., and Tin H.N. All authors reviewed the results and approved the final version of the manuscript.

LIST OF ABBREVIATIONS

AIDS	= Acquired Immunodeficiency Syndrome
COVID-19	= Coronavirus Disease 2019
HIV	= Human Immunodeficiency Virus
MSM	= Men who have sex with men
STDs	= Sexually Transmitted Diseases
WHO	= World Health Organization

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study was approved by the Ethics Committee in Biomedical Research at Can Tho University of Medicine and Pharmacy (No. 576/PCT-HDDD on July 15th, 2022).

HUMAN AND ANIMAL RIGHTS

The procedure performed in this study involving human participants was by the ethical standards of research committees and with the 1975 Declaration of Helsinki, as revised in 2013.

CONSENT FOR PUBLICATION

Informed consent was received from the participants.

STANDARDS OF REPORTING

STROBE guidelines were followed.

AVAILABILITY OF DATA AND MATERIALS

The authors confirm that the data supporting the findings of this research are available within the article. The corresponding author may share the collected data of the study upon reasonable request.

FUNDING

Declared none.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

The authors are grateful to Can Tho University of Medicine and Pharmacy in Vietnam for the time and effort they devoted to the study. They also thank Le Van Thinh Hospital for their support during the sample collection period.

SUPPLEMENTARY MATERIAL

Supplementary material is available on the publisher's website along with the published article.

REFERENCES

- [1] Low N, Broutet NJ. Sexually transmitted infections—research priorities for new challenges. CA, USA: Public Library of Science San Francisco 2017; p. e1002481.
- [2] Fasciana T. Sexually transmitted diseases: Diagnosis and control. Basel, Switzerland: MDPI 2022; p. 5293.
- [3] Agaçfidan A, Kohl P. Sexually transmitted diseases (STDs) in the world. FEMS Immunol Med Microbiol 1999; 24(4): 431-5. [http://dx.doi.org/10.1016/S0928-8244\(99\)00060-7](http://dx.doi.org/10.1016/S0928-8244(99)00060-7) PMID: 10435762
- [4] Wijesooriya NS, Rochat RW, Kamb ML, *et al.* Global burden of maternal and congenital syphilis in 2008 and 2012: A health systems modelling study. Lancet Glob Health 2016; 4(8): e525-33. [http://dx.doi.org/10.1016/S2214-109X\(16\)30135-8](http://dx.doi.org/10.1016/S2214-109X(16)30135-8) PMID: 27443780
- [5] Komar OM, Pidlisna IV. Analysis of the main indicators of socially dangerous sexually transmitted diseases. Rep Vinnyt Nat Med Univ 2020; 24(4): 709-13. [http://dx.doi.org/10.31393/reports-vnmedical-2020-24\(4\)-26](http://dx.doi.org/10.31393/reports-vnmedical-2020-24(4)-26)
- [6] Samkange-Zeeb FN, Spallek L, Zeeb H. Awareness and knowledge of sexually transmitted diseases (STDs) among school-going adolescents in Europe: A systematic review of published literature. BMC Public Health 2011; 11(1): 727. <http://dx.doi.org/10.1186/1471-2458-11-727> PMID: 21943100
- [7] Control, CfD and Prevention, Sexually transmitted disease surveillance 2018. Atlanta: US Department of Health and Human Services 2019; p. 10.
- [8] Ott MA, Santelli J. Sexually transmitted infections, public health, and ethics. The Oxford Handbook of Public Health Ethics. Oxfordshire, England: Oxford Academic 2019; p. 378.
- [9] Barrow RY, Ahmed F, Bolan GA, Workowski KA. Recommendations for providing quality sexually transmitted diseases clinical services, 2020. MMWR Recomm Rep 2020; 68(5): 1-20. <http://dx.doi.org/10.15585/mmwr.rr6805a1> PMID: 31899459
- [10] Rietmeijer CA. Improving care for sexually transmitted infections. J Int AIDS Soc 2019; 22(S6): e25349. <http://dx.doi.org/10.1002/jia2.25349> PMID: 31468743
- [11] Rhodes SD, Daniel-Ulloa J, Wright SS, *et al.* Critical elements of community engagement to address disparities and related social

- determinants of health: The centers of disease control and prevention community approaches to reducing sexually transmitted disease initiative. *Sex Transm Dis* 2021; 48(1): 49-55.
<http://dx.doi.org/10.1097/OLQ.0000000000001267> PMID: 32826480
- [12] Oluwale EO, Oyekanmi OD, Ogunyemi DO, Osanyin GE. Knowledge, attitude and preventive practices of sexually transmitted infections among unmarried youths in an urban community in Lagos State, Nigeria. *Afr J Prim Health Care Fam Med* 2020; 12(1): e1-7.
<http://dx.doi.org/10.4102/phcfm.v12i1.2221> PMID: 32370529
 - [13] Tenaw LA. *Bacterial Sexually Transmitted Disease*. London, England: IntechOpen 2022; pp. 1-6.
 - [14] Sexuality and migration: Milieu-specific access routes for sexual education of young people. *Results Represent Study Liv Environ* 2010; 14-7.
 - [15] Godeau E, Gabhainn NS, Vignes C, Ross J, Boyce W, Todd J. Contraceptive use by 15-year-old students at their last sexual intercourse: Results from 24 countries. *Arch Pediatr Adolesc Med* 2008; 162(1): 66-73.
<http://dx.doi.org/10.1001/archpediatrics.2007.8> PMID: 18180415
 - [16] Global health sector strategy on sexually transmitted infections 2016-2021. Geneva, Switzerland: World Health Organization 2016; pp. 1-8.
 - [17] Nguyen LTT, Tran CD, Nguyen HTT, *et al.* Nutritional status of Vietnamese infants assessed by Fenton growth chart and related factors: A cross-sectional study. *Pediatr Investig* 2023; 7(4): 254-62.
<http://dx.doi.org/10.1002/ped4.12394> PMID: 38050535
 - [18] Duong LM, Pham LV, Pham TT, *et al.* DNA-HPV transition rate and related factors in HPV-infected women in Can Tho city, Vietnam. *Trop Med Int Health* 2019; 24(11): 1330-4.
<http://dx.doi.org/10.1111/tmi.13309> PMID: 31520562
 - [19] Lan PT, Mogren I, Phuc HD, Lundborg CS. Knowledge and practice among healthcare providers in rural Vietnam regarding sexually transmitted infections. *Sex Transm Dis* 2009; 36(7): 452-8.
<http://dx.doi.org/10.1097/OLQ.0b013e31819fe9ae> PMID: 19556937
 - [20] Visalli G, Picerno I, Vita G, Spataro P, Bertuccio MP. Knowledge of sexually transmitted infections among younger subjects of the city of Messina (Sicily). *J Prev Med Hyg* 2014; 55(1): 17-22. PMID: 25916027
 - [21] Workowski KA, Bachmann LH, Chan PA, *et al.* Sexually transmitted infections treatment guidelines, 2021. *MMWR Recomm Rep* 2021; 70(4): 1-187.
<http://dx.doi.org/10.15585/mmwr.rr7004a1> PMID: 34292926
 - [22] Smith CJ. Social geography of sexually transmitted diseases in China: Exploring the role of migration and urbanisation. *Asia Pac Viewp* 2005; 46(1): 65-80.
<http://dx.doi.org/10.1111/j.1467-8373.2005.00260.x>
 - [23] Patterson-Lomba O, Goldstein E, Gómez-Liévano A, Castillo-Chavez C, Towers S. Per capita incidence of sexually transmitted infections increases systematically with urban population size: A cross-sectional study. *Sex Transm Infect* 2015; 91(8): 610-4.
<http://dx.doi.org/10.1136/sextrans-2014-051932> PMID: 25921021
 - [24] Alirol E, Getaz L, Stoll B, Chappuis F, Loutan L. Urbanisation and infectious diseases in a globalised world. *Lancet Infect Dis* 2011; 11(2): 131-41.
[http://dx.doi.org/10.1016/S1473-3099\(10\)70223-1](http://dx.doi.org/10.1016/S1473-3099(10)70223-1) PMID: 21272793
 - [25] Do SN. Predictive validity of the simplified Radiographic Assessment of Lung Edema score for the mortality in critically ill COVID-19 patients with the B. 1.617. 2 (Delta) variant in Vietnam: A single-centre, cross-sectional study. *medRxiv* 2024; 2024.03.
<http://dx.doi.org/10.1101/2024.03.20.24304599>
 - [26] Kreisel KM, Spicknall IH, Gargano JW, *et al.* Sexually transmitted infections among US women and men: Prevalence and incidence estimates, 2018. *Sex Transm Dis* 2021; 48(4): 208-14.
<http://dx.doi.org/10.1097/OLQ.0000000000001355> PMID: 33492089
 - [27] Tan PY, Somphos VS, Nguyen SD, Tan X. Demographic variation and socioeconomic inequalities associated with the triple burden of malnutrition in Vietnamese children aged 6 months to 9 years old: Findings from the Vietnamese General Nutrition Survey 2020. *medRxiv* 2024; 2024.03.
<http://dx.doi.org/10.1101/2024.03.18.24304456>
 - [28] Guidelines for the management of symptomatic sexually transmitted infections. Geneva, Switzerland: World Health Organization 2021; pp. 1-8.
 - [29] Mendiratta V, Meena AK, Verma D. Epidemiology and changing trends of sexually transmitted diseases over the past 17 years in a tertiary care center. *Indian J Sex Transm Dis AIDS* 2023; 44(2): 152-7.
http://dx.doi.org/10.4103/ijstd.ijstd_100_22 PMID: 38223154
 - [30] Nguyen SH, Dang AK, Vu GT, *et al.* Lack of knowledge about sexually transmitted diseases (STDs): Implications for STDs prevention and care among dermatology patients in an urban city in Vietnam. *Int J Environ Res Public Health* 2019; 16(6): 1080.
<http://dx.doi.org/10.3390/ijerph16061080> PMID: 30917565
 - [31] Adzrago D, Nyarko SH, Ananaba N. Effects of mental health and substance use disorder symptoms on the association between stds and sexual identity: Evidence from the 2017-2019 national survey on drug use and health. *J Public Health* 2021; 31: 1225-36.
<http://dx.doi.org/10.1007/s10389-021-01683-y>
 - [32] Tan NV, Kien NT. Assessment of access to interventions to prevent and control sexually transmitted diseases in men who have sex with men in Ben Tre province period 2014 - 2018. *Can Tho J Med Pharma* 2020; (27): 126-34.
 - [33] Munain DJL. Epidemiology and current control of sexually transmitted infections. The role of STI clinics. *Enferm Infecc Microbiol Clin* 2019; 37(1): 45-9.
<http://dx.doi.org/10.1016/j.eimc.2018.10.015>
 - [34] Tran GV, Hayashida T, Dang ALD, *et al.* Prevalence of transmitted drug resistance and phylogenetic analysis of HIV-1 among antiretroviral therapy-naïve patients in Northern Vietnam from 2019 to 2022. *Global Health Med* 2024; 6(2): 117-23.
<http://dx.doi.org/10.35772/ghm.2023.01112> PMID: 38690128
 - [35] Brookmeyer KA, Haderxhanaj LT, Hogben M, Leichter J. Sexual risk behaviors and STDs among persons who inject drugs: A national study. *Prev Med* 2019; 126: 105779.
<http://dx.doi.org/10.1016/j.ypmed.2019.105779> PMID: 31319117
 - [36] Hua TT, Nguyen TH, Nguyen TH, Thi Hue Cao C, Nguyen TH. Roles and current challenges of peer educators in delivering hiv/aids-related services: A case study in vietnam. *F1000 Res* 2024; 13: 707.
<http://dx.doi.org/10.12688/f1000research.152867.1>
 - [37] Thu HNT, Hung PT, Ha TNT, *et al.* Restructure of the provincial preventive medicine system in Vietnam: Implication for the HIV/AIDS human resources. *Int J Healthc Manag* 2024; 17(4): 734-42.
<http://dx.doi.org/10.1080/20479700.2023.2234142>
 - [38] Tuan AN, Fylkesnes K, Thang BD, *et al.* Human immunodeficiency virus (HIV) infection patterns and risk behaviours in different population groups and provinces in Viet Nam. *Bull World Health Organ* 2007; 85(1): 35-41.
<http://dx.doi.org/10.2471/BLT.05.026153> PMID: 17242756
 - [39] Global progress report on HIV, viral hepatitis and sexually transmitted infections. Geneva, Switzerland: World Health Organization 2021; pp. 1-112.
 - [40] Newby KV, Wallace LM, French DP. How do young adults perceive the risk of chlamydia infection? A qualitative study. *Br J Health Psychol* 2012; 17(1): 144-54.
<http://dx.doi.org/10.1111/j.2044-8287.2011.02027.x> PMID: 22233107
 - [41] Paganella MP, Motta DLR, Adami ADG, Sperhake RD, Kato SK, Pereira GFM. Knowledge about sexually transmitted infections among young men presenting to the Brazilian Army, 2016. *Medicine* 2021; 100(22): e26060.
<http://dx.doi.org/10.1097/MD.00000000000026060> PMID: 33492089

- 34087848
- [42] Lan PT, Nguyen HT, Golparian D, Van NTT. The WHO enhanced gonococcal antimicrobial surveillance programme (EGASP) identifies high levels of ceftriaxone resistance across vietnam. *Lancet Reg Health West Pacif* 2023; 48: 101125. <http://dx.doi.org/10.1016/j.lanwpc.2024.101125> PMID: 39040039
- [43] Mansor N, Ahmad N, Rahman HA. Determinants of knowledge on sexually transmitted infections among students in public higher education institutions in Melaka state, Malaysia. *PLoS One* 2020; 15(10): e0240842. <http://dx.doi.org/10.1371/journal.pone.0240842> PMID: 33119620
- [44] Tran VC, Nguyen SH, Bui HT. Impact of gender, location, and seasonality on the prevalence of gonorrhoea and chlamydia co-infections in northern vietnam. *Indian J Microbiol* 2024; 1-9. <http://dx.doi.org/10.1007/s12088-024-01301-x>
- [45] Nguyen H, Ngoc DA, Le NV, *et al.* Prevalence, risk factors and genotyping of *chlamydia trachomatis* from endocervical specimens of infertile women at a tertiary care hospital, Vietnam. *Int J STD AIDS* 2024; 35(6): 452-61. <http://dx.doi.org/10.1177/09564624241230342> PMID: 38294256
- [46] Nguyen PTT, Pham HV, Van DH, Pham LV, Nguyen HT, Nguyen HV. Randomized controlled trial of the relative efficacy of high-dose intravenous ceftriaxone and oral cefixime combined with doxycycline for the treatment of Chlamydia trachomatis and Neisseria gonorrhoeae co-infection. *BMC Infect Dis* 2022; 22(1): 607. <http://dx.doi.org/10.1186/s12879-022-07595-w> PMID: 35810277
- [47] Vanessa G A. Study raises concern over cefixime failure rate for gonorrhoea. 2013. Available from: <https://www.managedhealthcareexecutive.com/view/study-raises-concern-over-cefixime-failure-rate-gonorrhea>
- [48] Nguyen LH, Le TBT, Le NQN, Tran NTT. Acceptance and willingness to pay for a vaccine against human papilloma virus (HPV) among parents of boys in central Vietnam. *Front Public Health* 2022; 10: 801984. <http://dx.doi.org/10.3389/fpubh.2022.801984> PMID: 35356024
- [49] Dombrowski JC, Wierzbicki MR, Newman LM, *et al.* Doxycycline versus azithromycin for the treatment of rectal Chlamydia in men who have sex with men: A randomized controlled trial. *Clin Infect Dis* 2021; 73(5): 824-31. <http://dx.doi.org/10.1093/cid/ciab153> PMID: 33606009
- [50] Lin X, Chen W, Yu Y, *et al.* Emergence and genomic characterization of Neisseria gonorrhoeae isolates with high levels of ceftriaxone and azithromycin resistance in Guangdong, China, from 2016 to 2019. *Microbiol Spectr* 2022; 10(6): e01570-22. <http://dx.doi.org/10.1128/spectrum.01570-22> PMID: 36377922
- [51] Adamson PC, Bhatia R, Tran KDC, *et al.* Prevalence, anatomic distribution, and correlates of Chlamydia trachomatis and Neisseria gonorrhoeae infections among a cohort of men who have sex with men in Hanoi, Vietnam. *Sex Transm Dis* 2022; 49(7): 504-10. <http://dx.doi.org/10.1097/OLQ.0000000000001626> PMID: 35312657
- [52] Nguyen TV, Tran HP, Khuu NV, *et al.* Increases in both HIV and syphilis among men who have sex with men in Vietnam: Urgent need for comprehensive responses. *Int J STD AIDS* 2021; 32(14): 1298-307. <http://dx.doi.org/10.1177/09564624211036421> PMID: 34392717

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.